


OPTOMETRIC EDUCATION

The Journal of the Association of Schools and Colleges of Optometry

Volume 45, Number 1
Fall 2019



Does Practice Make Perfect? Relating Student Performance to Training Hours

Student Performance and Attitudes During the Transition from Paper to Computer-Based Examinations

Development of a Polymeric Eye Model for Foreign Body Removal

Optic Nerve Melanocytoma: a Teaching Case Report

Lymphoproliferative Disorders of the Ocular Adnexa: Teaching Case Reports of Conjunctival MALT Lymphoma and Lymphoid Hyperplasia

ALSO INSIDE

Editorial: The Use of Social Media to Enhance Academic Careers

ASCO's Public Awareness Campaign "Optometry Gives Me Life" Surpasses Benchmarks

Announcement: *Optometric Education* Welcomes Associate Editors

Optometric Education: Volume 45, Number 1 (Fall 2019)

Table of Contents

Articles

[Does Practice Make Perfect? Relating Student Performance to Training Hours](#)

[Student Performance and Attitudes During the Transition from Paper to Computer-Based Examinations](#)

[Development of a Polymeric Eye Model
for Foreign Body Removal](#)

[Lymphoproliferative Disorders of the Ocular Adnexa: a Teaching Case Report of Conjunctival MALT Lymphoma and Lymphoid Hyperplasia](#)

Features

[The Use of Social Media to Enhance Academic Careers](#)

[ASCO's Public Awareness Campaign "Optometry Gives Me Life" Surpasses Benchmarks](#)

[Two Join ASCO Journal as Associate Editors](#)

Articles

PEER REVIEWED

Does Practice Make Perfect? Relating Student Performance to Training Hours

Yutaka Maki, OD, MS, FCOVD, and Brian K. Foutch, OD, PhD, FAAO | *Optometric Education: Volume 45, Number 1 (Fall 2019)*

Abstract

We tested the hypothesis that a positive correlation exists between time practiced outside of scheduled laboratories and high-stakes proficiency performance in a preclinical optometry course. Practice hours for 130 first-year optometry students were recorded weekly and compared with the average scores of two summative ? midterm and final ? proficiency examinations. Due to scheduling constraints, the final proficiency test was administered during weeks 12 and 14, and students were randomly assigned to either week. The group scheduled for week 14 practiced significantly more hours (85.7 ± 19.4 hours vs. 73.0 ± 16.8 hours) and, on average, performed significantly better ($89.7 \pm 6.1\%$ vs. $85.9 \pm 7.3\%$) than the week-12 group. At the individual level within each group, however, we found a negative correlation between practice hours and performance, particularly for the week-14 group. This study showed that students did not necessarily perform better by practicing more. The results are discussed in the context of clinical learning aspects such as deliberate practice and self-awareness of current skill set.

Key Words: assessments, preclinical, laboratory, practice, self-awareness

Background

We have all heard the phrase “practice makes perfect.” The Latin proverb “usus est magister optimus” translates to “practice is the best master,” and Aristotle said, “For the things we have to learn before we can do them, we learn by doing them.”¹ Emphasis on the importance of practice is seen clearly in every culture and century, and it seems safe to assume that the more one practices, the better one performs. If this assumption were true, it would make sense to assign students struggling in preclinical optometry laboratory courses additional practice outside of scheduled laboratory times. By extension, it would be reasonable to assume that additional practice should result in higher grades or evaluations on preclinical assessments. In this study, we examined whether students who practiced more hours outside of their assigned course laboratory performed better on high-stakes midterm and final proficiency assessments.

Methods

We retrospectively reviewed the course records of 130 first-year optometry students enrolled in the second semester preclinical course during Spring 2015 and 2016. The data were collected in the clinical laboratory facilities at the University of the Incarnate Word Rosenberg School of Optometry (UIWRSO), San Antonio, Texas, under the supervision of the course instructors.



Table 1. [Click to enlarge](#)

The students learned new clinical skills (retinoscopy, refraction, binocular tests) each week in the three-hour course laboratory (**Table 1**). To ensure they were reviewing the newly learned skills, all students were required to practice a minimum of four hours in the laboratory outside of instruction time through week 10. The newly learned skills were evaluated in the form of checkouts (short clinical skill assessments) throughout the semester (Table 1). The primary purpose of the checkouts was to provide students with formative feedback to learn from their successes and mistakes. A secondary goal was for students to practice more if they performed poorly. To ensure this was taking place, two additional hours were mandated for each checkout with a grade below 85%. Thus, students who passed or failed a checkout were required to practice a minimum of four or six hours, respectively. Beyond that, no additional assignments were given. However, all students were encouraged to practice more than the minimum assigned hours, especially if they felt they had not reached expected skill levels.

When students practiced outside of the scheduled laboratory, they were under the supervision of teaching assistants. Each student recorded the time he or she entered and exited the laboratory, and the teaching assistant confirmed the times. Each week's logged hours were then submitted to the instructor of record in the subsequent course laboratory. The three scheduled hours they spent each week in the course laboratory were not included in the logged hours.

In addition to the checkouts, there was one midterm proficiency assessment (retinoscopy and refraction) and one final proficiency assessment (entire refractive sequence of lensometry, entrance tests, retinoscopy, subjective refraction and photometry). The rubrics used in checkouts and proficiency assessments were modeled after the Clinical Skills Exam evaluation forms used by the National Board of Examiners in Optometry.² Unlike weekly checkouts, no additional practice hours were assigned to students who scored below 85% for the proficiency assessments.

We recorded the number of hours each student practiced up to their scheduled proficiency assessments. Because the final assessments were administered during weeks 12 and 14 of the semester, we randomly divided the students into two groups for analysis. The students' cumulative grades from the midterm (33%) and final (67%) proficiency assessments were compared with the number of hours they practiced to determine whether there was any correlation (Pearson's r reported). Checkout grades were excluded from the cumulative grade for the purpose of this retrospective analysis because the mandated additional two-hour assignment for failing checkouts inevitably would have decreased any positive relationship between the grade and the practice hours. In our analyses, we defined statistically significant as p -value less than 0.05.

Results

Overall, the second (week-14) group had higher mean cumulative lab grades and mean number of logged practice hours ($89.7 \pm 6.1\%$, 85.7 ± 19.4 hours) than the first (week-12) group ($85.9 \pm 7.3\%$, 73.0 ± 16.8 hours). The differences were statistically significant ($p < 0.01$) for both the cumulative lab grades and the logged practice hours (**Tables 2 and 3**).

However, we found a negative correlation between cumulative grades and overall practice hours within each group. Within each group, the finding was statistically significant for the week-14 group ($r = -0.30$, $p = 0.02$) (**Figure 1 and Table 4**) but not for the week-12 group ($r = -0.02$, $p = 0.84$) (**Figure 2 and Table 4**). Similar patterns were seen when comparing final proficiency grades with the *overall* practice time (Table 4) as well as the cumulative grades with *voluntary* practice time (**Table 5**).



Figure 1. [Click to enlarge](#)



Figure 2. [Click to enlarge](#)



Table 2. [Click to enlarge](#)



Table 3. [Click to enlarge](#)



Table 4. [Click to enlarge](#)



Table 5. [Click to enlarge](#)

There was no statistically significant difference in practice time between the week-12 and week-14 groups up to the midterm proficiency assessment or the last (week 10) checkout ($p=0.718$ and $p=0.487$ respectively, Table 2). Also, there was no statistically significant difference in checkout and midterm grades between the week-12 and week-14 groups ($p=0.473$ and $p=0.052$ respectively, Table 3).

Further analysis of the correlation between post-midterm practice time and midterm proficiency grades revealed a statistically significant correlation for the week-14 group but not the week-12 group ($p<0.01$ and $p=0.25$ respectively) (**Table 6**).

When correlations among each assessment grade were analyzed, the final proficiency grades for the week-12 group were significantly positively correlated with midterm proficiency and checkout grades ($p<0.01$ each). This was not the case for the week-14 group ($p=0.42$ and $p=0.05$ for midterm proficiency and checkout grades respectively) (**Table 7**).



Table 6. [Click to enlarge](#)



Table 7. [Click to enlarge](#)

Discussion

The study results supported the conventional wisdom: “practice makes perfect.” The group given an additional two weeks increased their practice time and performed better. Although it could be argued that these results can be explained by the early group giving the latter group a “heads-up” on the proficiency assessments, we believe it would have given them little to no advantage. The final proficiency assessment was comprised of elements from all the checkouts, so there were few surprises. In addition, the same final proficiency assessment was given in previous years; therefore, students from either group could have asked upperclassmen about it.

At the individual level, we found different results within each group. Higher-performing students in the week-14 group practiced less than lower-performing students. This was not the case for the week-12 group, where we found no statistically significant correlation between practice and performance.

Many factors could explain this apparent contradiction in the week-14 group. First, students have different starting skill levels based on work or academic experience. Second, some learn new skills more innately than others, and we have informally observed that it is often the struggling students who practice more to compensate for their current lack of skills. In fact, the statistically significant negative correlation between post-midterm practice time and midterm proficiency grades of the week-14 group indicates that the outcome of the midterm proficiency assessment affected their practice behavior (Table 6). While we had anticipated that the effort shown by their increased practice time would be the predictor for their clinical performance, we have instead found their current clinical progress to be the predictor of their effort. That is, lower-performing students practiced more.

This type of self-adjusted practice behavior could be explained by emotional intelligence (EI), which is defined as the ability to monitor one's own and others' emotions, to discriminate among them, and to use this information to guide one's thinking and actions.³ One model of EI proposed by Goleman and Boyatzis identifies four clusters of competencies: self-awareness, self-management, social awareness and relationship management.⁴ All of these clusters appear to be important predictors of clinical success, and several authors have observed positive associations between higher EI and better clinical performance in dental, medical and nursing school students.⁵⁻⁹ In one of the studies, self-management competencies were significantly correlated with student clinical performance (as measured by mean clinical grade).⁵ A recent study of optometry students in the United Kingdom demonstrated a positive association between self-awareness and academic performance.¹⁰ The characteristics of self-management include adaptability, initiative, persistence in pursuing goals, taking responsibility for personal performance, and striving to meet a standard of excellence.¹¹ This also may explain the practice pattern of our higher-performing students. Perhaps they based their decision to practice less on either positive feedback received during weekly checkouts, proper time management of overall curriculum demands, or awareness of their clinic skills.

While considering EI may offer insight into the week-14 group's negative correlation between practice and performance, we observed no statistically significant correlations between assessment grades (midterm proficiency, final proficiency or cumulative) and practice time (Tables 4-6) in the week-12 group. Because the major difference between the two groups was the amount of practice time given, this suggests availability of time is a key factor in behavioral change in students. This also explains why statistically significant correlations appeared between checkouts and proficiency assessments (midterm or final) in the week-12 group, while no such correlations appeared in the week-14 group (Table 7). Because the week-12 group lacked sufficient time to adapt their learning and close the gap in their achievement, midterm proficiency performance became the predictor for their final proficiency result.

This suggests the importance of making struggling students fully aware of their initial (or current) status and giving adequate support and time to close any achievement gap. Although we initially believed that weekly checkouts and the midterm proficiency assessment would be sufficient to provide such awareness, some of the lower-performing students did not respond to this feedback as we expected. When analyzing the bottom 20% of the week-12 and week-14 groups, we found that 57.1% and 33.3% of students were practicing below the average practice time, respectively. In the week-12 group, the average group and bottom 20% practice times were 73.0 ± 16.8 hours and 72.9 ± 16.1 hours, respectively. In the week-14 group, the average group and bottom 20% practice times were 85.7 ± 19.4 hours and 93.5 ± 20.3 hours, respectively. This indicates that identifying and guiding low-performing students who lack self-awareness and/or self-management earlier in the course is essential. Convincing students of their status might be challenging, however, as at least two previous investigations of competence awareness have shown that lower-achieving students overestimate their competence and vice versa for higher-achieving students (an effect that held even after feedback was given, though to a lesser extent).^{12,13}

When offering support to low-performing students, it is important to teach them how to practice. Studies

show that expert performance is most effectively attained by deliberate practice (DP), where focused training is designed and arranged by teachers and coaches to optimize improvement.¹⁴⁻¹⁶ This focused training also involves the provision of immediate feedback, time for problem-solving and evaluation, and opportunities for repeated performance to refine behavior. In a preclinical lab setting, perhaps the lower-performing students can be given individualized assignments based on their current performance and/or be paired with a teaching assistant or a high-performing student to focus on specific areas of struggle.

In the current study we did not provide any designed training, nor did we measure how each student actively engaged in DP. This certainly limits any inferences made about the effect of DP on performance. Further limitations of our study included a relatively small sample size and a retrospective period of only two years. Lastly, our study lacked any direct measures of EI or self-awareness. Further studies could and should do so by integrating focus group interviews with higher- and lower- achieving students and teaching assistants into the analysis.

Conclusion

Our study showed that individual performance depended on amount of practice and feedback from assessments. Although at the class level the group that had more time to practice performed better, at the individual level within each group students who practiced more did not necessarily perform better. Our findings suggested that students adjusted the number of hours they practiced according to their perception of their current skill set or a self-prescribed mastery goal. Ultimately, assessment performance and students' practice behavior influenced each other. In addition, we found it to be essential to provide sufficient time for students to adapt and reach the competency level, otherwise earlier assessment results become the predictor for their final outcome. Lastly, it is imperative to make struggling students aware of their status and provide enough support as early as possible. While our findings have implications for medical and optometric education, they will be strengthened by more direct measurements of DP, self-awareness and EI. As a result of this analysis, we have considered providing all students at least three weeks to practice before major assessments. In addition, underperforming students will be prescribed a deliberate practice plan and be scheduled in the later assessment week.

Acknowledgements

A portion of this work was presented at the 2016 annual meeting of the American Academy of Optometry in Anaheim, Calif. The authors acknowledge the University of the Incarnate Word Rosenberg School of Optometry classes of 2018 and 2019 who served as subjects of this study.

References

1. Broadie S. Ethics with Aristotle. Oxford University Press, Incorporated. 1991.
2. National Board of Examiners in Optometry – NBEO [Internet]. Candidate Eligibility – NBEO. [cited 2017 Dec 17]. Available from: <https://www.optometry.org/nccto.cfm>
3. Mayer JD, Salovey P. What is emotional intelligence? In: Salovey P, Sluyter DJ, eds. Emotional development and emotional intelligence: educational implications. New York, NY: Basic Books 1997;3-31.
4. Goleman D, Boyatzis RE, McKee A. Primal leadership: realizing the power of emotional intelligence. Boston: Harvard Business School Press, 2002.
5. Victoroff KZ, Boyatzis RE. What is the relationship between emotional intelligence and dental student clinical performance? J Dent Educ. 2013 Apr;77(4):416-26.
6. Beauvais AM, Brady N, O'Shea ER, Griffin MT. Emotional intelligence and nursing performance among nursing students. Nurse Educ Today. 2011 May;31(4):396-401.
7. Arora S, Ashrafian H, Davis R, Athanasiou T, Darzi A, Sevdalis N. Emotional intelligence in medicine: a systematic review through the context of the ACGME competencies. Med Educ. 2010

Aug;44(8):749-64.

8. Satterfield J, Swenson S, Rabow M. Emotional intelligence in internal medicine residents: educational implications for clinical performance and burnout. *Ann Behav Sci Med Educ.* 2009;14(2):65-68.
9. Stratton TD, Elam CL, Murphy-Spencer AE, Quinlivan SL. Emotional intelligence and clinical skills: preliminary results from a comprehensive clinical performance examination. *Acad Med.* 2005 Oct;80(10 Suppl):S34-7.
10. Pancholi BR, Dunne MCM. Virtual patient instruction and self-assessment accuracy in optometry students. *Optometric Education.* 2018;43(2):1-16.
11. Goleman D. Emotional intelligence. April 21, 2015. Accessed: October 5, 2017. Available from: www.danielgoleman.info/daniel-goleman-how-emotionally-intelligent-are-you/.
12. Hodges B, Regehr G, Martin D. Difficulties in recognizing one's own incompetence: novice physicians who are unskilled and unaware of it. *Acad Med.* 2001 Oct;76(10 Suppl):S87-9.
13. Kruger J, Dunning D. Unskilled and unaware of it: how difficulties in recognizing one's own incompetence lead to inflated self-assessments. *J Pers Soc Psychol.* 1999 Dec;77(6):1121-34.
14. Ericsson KA. Deliberate practice and acquisition of expert performance: a general overview. *Acad Emerg Med.* 2008 Nov;15(11):988-94.
15. McGaghie WC, Issenberg SB, Cohen ER, Barsuk JH, Wayne DB. Does simulation-based medical education with deliberate practice yield better results than traditional clinical education? A meta-analytic comparative review of the evidence. *Acad Med.* 2011 Jun;86(6):706-11.
16. Plant EA, Ericsson KA, Hill L, Asberg K. Why study time does not predict grade point average across college students: implications of deliberate practice for academic performance. *Contemporary Educational Psychology.* 2005;30(1):96-116.

Yutaka Maki, OD, MS, FCOVD, [maki@uiwtx.edu] is an Assistant Clinical Professor at the University of the Incarnate Word (UIW) Rosenberg School of Optometry and Chief of the Binocular Vision and Vision Therapy Service at UIW Eye Institute. His didactic responsibilities include teaching courses in optics, preclinical optometry, vision therapy and strabismus/amblyopia.

Brian K. Foutch, OD, PhD, FAAO, is an Associate Professor at the University of the Incarnate Word Rosenberg School of Optometry where he teaches public health and epidemiology and neuroanatomy.

PEER REVIEWED

Student Performance and Attitudes During the Transition from Paper to Computer-Based Examinations

Brian K. Foutch, OD, PhD, Lourdes Fortepiani, MD, PhD, and Richard Trevino, OD | *Optometric Education: Volume 45, Number 1 (Fall 2019)*

Abstract

Purpose: To compare paper-based test (PBT) and computer-based test (CBT) scores and whether demographics, previous academic success, learning style or attitudes affected performance. **Methods:** 65 second-year optometry students completed attitude questionnaires after four midterm examinations — one PBT and one CBT in two courses taught by the same instructor in the same semester. Attitudes were analyzed via χ^2 comparisons while test scores were compared via analysis of variance. **Results:** PBT and CBT scores were the same, but female, non-Hispanic and non-White students and those with a preference for PBT had higher PBT scores. **Conclusion:** While overall PBT and CBT formats appeared equivalent, we uncovered demographic trends in both test performance and attitudes that warrant further investigation.

Key Words: computer-based test, paper-based test, student performance

Background

Standardized assessments used for professional licensure and to measure pre-professional aptitude have been administered electronically for decades. Computer-based test (CBT) platforms have also been introduced in many undergraduate and professional programs, and their use has coincided with the emergence of cognitive learning theories that stress integration of classroom teaching and assessment.¹

Most health profession programs have multiple layers of learning outcomes. Specifically, optometric educators must consider individual course objectives, school or overall curriculum outcomes, and professional outcomes and attributes defined by the Association of Schools and Colleges of Optometry (ASCO),² the Accreditation Council on Optometric Education (ACOE)³ and the National Board of Examiners in Optometry (NBEO).⁴ While the 20th century paradigm of providing traditional lectures followed by static, multiple-choice paper-based tests (PBT) still exists and has value,⁵ it makes it far more challenging and time-consuming to measure student performance on these learning outcomes across a curriculum. After all, PBT formats do not automatically generate reports of student performance according to learning outcomes in a course or across the curriculum. That process requires a significant time investment and coordination between instructors after a test is given. In contrast, CBT platforms allow evaluators to digitally assign learning levels and course or program levels to test questions, providing improved feedback when attempting to drive assessment-based curriculum or instructional changes.⁶ Further, controlled studies and informal observations have shown that CBT formats provide additional advantages as well as some disadvantages for students and instructors. Additional advantages include automatic grading with nearly real-time score reporting to students⁷ and higher-resolution feedback about specific subject content.⁸ Disadvantages of CBT formats include computer eyestrain, compatibility and connectivity issues⁸ as well as difficulties creating, navigating or grading in-depth questions.⁹ Arguably, the most important barrier is the general skepticism among both instructors

and students toward computerized testing until they achieve an initial positive experience.¹⁰

Previous studies of student academic performance using CBT are equivocal. At least two groups of researchers have concluded that simply administering an exam using a computerized format vs. a traditional paper- and pencil-based format had no significant effect on achievement^{11,12} while another group has observed that test scores improved with CBT formats for both static-type (multiple-choice, true/false) and interactive problem-based questions.¹³ Our institution recently implemented one such CBT platform, and computerized testing was highly encouraged (but optional) by our administration during a “hybrid” period of one academic year. At the end of that year, instructors were required to administer only computerized tests. That one-year period ? when both PBT and CBT formats were used ? provided us with a unique opportunity to better understand how student demographics, academic performance and attitudes influenced their transition to computerized testing. Our aim was to determine whether there was a significant difference between mean PBT and CBT scores and the extent to which various student factors and attitudes could influence performance. The results from this study could reveal different strategies for improving student performance and easing the overall transition to computer-based testing.

Methods

Subjects

We recruited 65 (39 females, 26 males) subjects with a mean age (\pm S.D.) of 24.7 (\pm 1.87) years. Due to the high proportion of Hispanics in our student population, we wanted to analyze the results not only considering age, gender and race but also Hispanic ethnicity (10 Hispanic, 55 non-Hispanic). Volunteers were eligible if they were enrolled and in good academic standing (i.e., cumulative GPA of 2.0 or higher) in the Rosenberg School of Optometry’s second-year class during the fall semester of 2015. All subjects were recruited via a classroom announcement by the investigators and were all paid for their participation. All subjects provided informed consent. The study protocol was approved by the institutional review board at the University of the Incarnate Word and carried out according to the guidelines set out in the 1964 Declaration of Helsinki.

Research design

During the fall semester of their second year, 65 students completed two midterm examinations in two different courses (ocular physiology and organ pathology). These courses were selected because they were offered simultaneously in the same semester by the same instructor, and all exams were delivered in the same classroom setting. This could minimize the impact of different instructors using different question styles to assess student performance. These examinations were administered as regularly scheduled assessments in each course. In the first course (organ pathology), the students took the first midterm electronically via ExamSoft (ExamSoft Worldwide, Inc., Boca Raton, Fla.) and the second midterm using a traditional paper and pencil format (graded using Scantron forms and software; Scantron Corporation, Eagan, Minn.). The order was reversed for the second course (ocular physiology). The overall crossover design is shown in **Figure 1**. Despite the possible influence of test performance on student attitudes, students were unaware of this research opportunity until after the end of the second midterms. This was done for two reasons. We were primarily interested in student performance and attitudes after their initial exposure to CBT. Students in this cohort were not previously exposed to CBT (ExamSoft) in optometry school but rather used PBT (Scantron) during their first year. In addition, we did not want to create anxiety nor prompt concerns in students about the new CBT format.



Figure 1. Subjects took two midterms in two separate courses (organ pathology and ocular physiology). In organ pathology,

the first midterm was administered electronically and the second midterm using a traditional paper and pencil format. The order was reversed for the second course. After both midterms, attitudes toward the examination formats were measured via a custom questionnaire.

[Click to enlarge](#)

While all examinations contained only multiple-choice items (both single and multiple response), the computerized format allowed for randomized ordering of questions and answer choices (i.e., multiple exam versions) while Scantron-graded exams were administered using a single version. Highlighting and backward navigation were the only additional features enabled on ExamSoft. In addition, students had access to scratch paper regardless of the testing platform. At the conclusion of the semester, study participants' attitudes toward paper and computerized exams were measured via a custom questionnaire with 5-rating scale semantic differential items (use and analysis described in a review of best practices in assessing student attitudes¹⁴).

With our semantic differential items, respondents did not indicate a level of agreement with a statement (as in Likert scaled items); rather, they chose a position on a scaled line that connoted a preference for one method or the other. For example, survey question #1 asked subjects to indicate whether they were "...more stressed taking the test using the computer-based test" or "...more stressed taking the test on paper." Whether the direction (+ or -) corresponded to CBT or PBT preference was randomly assigned for each question. However, for analysis, "-2" (on the 5-rating scale) indicated a strong preference for the PBT (more stressed by the CBT) and a "+2" indicated a strong preference for CBT (more stressed by the PBT). Responses of "-1" or "+1" indicated less endorsement, and a "0" indicated no preference or endorsement at all. Students also completed a 16-item VARK questionnaire (VARK; available online at <https://vark-learn.com/the-vark-questionnaire/>) to determine the level to which they are visual (V), aural (A), read/write (R) or kinesthetic (K) learners, as different learning styles have been associated with student performance.¹⁵

Analysis

Our first goal was to determine the relationship between the CBT and PBT scores. We examined scores across the four examinations via repeated measures analysis of variance (ANOVA) with course and examination mode as fixed factors. We then pooled all the exam scores into a single variable representing each subject's difference between their average CBT and PBT scores. These differences were also analyzed by ANOVA with gender, race (White vs. non-White) and ethnicity (Hispanic vs. non-Hispanic) as fixed factors. The difference scores were also regressed on continuous variables such as age and academic parameters (Optometry Admission Test [OAT] scores, undergraduate grade point averages [GPAs], GPA thus far in the optometry program, and learning style and attitudes). Distributions of subjects by demographics are shown in **Table 1**.



Table 1. [Click to enlarge](#)

We determined the level of endorsement (either for the CBT or PBT format) by comparing the medians for each survey question to zero via a Wilcoxon signed rank test. Ad hoc comparisons of students performing best on PBT or CBT were also performed, and these are described in detail in the Results section. Statistical significance was defined as a p-value less than 0.05 for all testing. All analyses were performed using SPSS (IBM Corporation, Armonk, N.Y.) and Excel (Microsoft Inc., Redmond, Wash.).

Results

All 65 second-year students consented to participation. The range of the difference between the

computer-based and paper-based scores (C-P, for convenience) was -18.6 to 13.9 (refer to **Table 2** for summary of scores). One male subject scored 34.5 points (4.5 standard deviations from the mean difference) better on the PBT. His performance and survey data were excluded from all analyses, as his performance bias for paper-based testing was considered an outlier. All test scores for the remaining subjects ($n = 64$) were analyzed using repeated measures ANOVA. The main effect of course [$F(1,57) = 11.43, p = 0.001$] — with students scoring 3.4% higher in ocular physiology — and the interaction of course and exam mode [$F(1,57)=6.607, p=0.013$] were both significant, but there was no main effect of exam mode [$F(1,57)=1.740, p=0.192$]. So, while there was a bias (C-P=-2.30) in our subjects toward better performance on the PBT, the difference was insignificant.



Table 2. [Click to enlarge](#)

For the two formats to be considered equivalent, it has been suggested that the two testing methods must yield similar dispersions and overall distributions of scores.¹⁶ The scores from all examinations were distributed normally ($p>0.05$ on Anderson Darling testing), though the distributions were shaped somewhat differently (**Figure 2**). The distribution most distinct from the others was for the ocular physiology PBT, where no student received a failing grade (<65%). However, 4.7% (3 of 64) of students failed the CBT in ocular physiology. This trend was reversed for organ pathology, where only 6.3% (4 of 64) failed the CBT, but 10.9% (7 of 64) failed the PBT. There does not appear then to be a systematic difference in test scores based on test platform. We then combined the two course test scores for each platform, and the relationship between the average scores from each platform is shown in **Figure 3**. Computer-based and paper-based scores were highly positively correlated ($r=0.62, p<0.001$), providing additional evidence that the two platforms are essentially equivalent in assessing student performance.



Figure 2. Distributions of paper-based test (PBT) and computer-based test (CBT) scores. The shapes of the distributions were somewhat different among all four midterm examinations, but all were distributed normally. [Click to enlarge](#)



Figure 3. Computer-based test scores and paper-based test scores were highly positively correlated. [Click to enlarge](#)

It has been argued, however, that any two methods that test the same parameter and provide wide variability in scores will almost always be positively correlated.¹⁷ It is better for us to calculate the difference (C-P) and plot this difference against the overall average of both methods.¹⁸ This difference (or Bland-Altman) plot is shown in **Figure 4**. The mean difference (-2.30) is represented by the solid line. When the difference between the two methods was regressed on the average of all exams, we saw no relationship ($r=0.06, p>0.9$). That is, mean overall test performance did not systematically predict a performance bias for either testing platform. The differences were distributed normally (Kolmogorov-Smirnov statistic=0.10, $p>0.05$) with a median of -3.50 and 25th and 75th percentiles of -6.63 and 2.25, respectively. This slightly positively skewed distribution was seen when we computed the reference interval (or 95% limit of agreement) for the two testing methods (mean ± 1.96 standard deviation of the difference). These values (10.3, -15.3) are shown in Figure 4 as thick dashed lines and indicated significant agreement between the two methods for all but six subjects (four performing significantly better on the CBT and two significantly better on the PBT).



Figure 4. Positive differences represent better mean performance on the CBT. The solid black line represents the mean difference between CBT and PBT scores (-2.30). The dotted lines represent the 25th percentile (-6.63) and 75th percentile (2.25). The heavy dashed lines represent the 95% confidence intervals below (mean -1.96 S.E.; -15.27) and above the mean (mean +1.96 S.E.; 10.57). [Click to enlarge](#)

While the testing administration methods appeared to be equivalent, we were still interested in what factors may be responsible for the trend in our subjects to perform better on paper. We pooled all the exam scores into a single variable (C-P), which was analyzed by ANOVA with gender, race (White vs. non-White) and ethnicity (Hispanic vs. non-Hispanic) as fixed factors. We found a trend for the mean bias towards PBT to be more in non-White subjects than in White subjects [$F(1,57)=3.093$, $p=0.053$]. When analyzed separately by one-sample t-tests, the bias toward PBT in non-White subjects was significantly less than zero [$t(57)=-3.22$, $p<0.005$] but not significant [$t(57)=-0.98$, $p=0.33$] in White subjects. The effects of gender [$F(1,57)=0.609$, $p=0.439$] and ethnicity [$F(1,57)=0.272$, $p=0.604$] were not significant. However, while neither gender nor ethnicity played a significant role on ANOVA, there was a difference when analyzed separately. That is, the mean difference for female subjects (-2.73) was significantly less than zero on a one-sample t-test [$t(57)=-2.75$, $p=0.01$]. This was not the case for male subjects [mean: -1.62; $t(57)=-1.11$, $p=0.28$]. Similarly, the mean for non-Hispanic subjects (-2.38) was significantly less than zero [$t(57)=-2.65$, $p=0.01$] but not for Hispanic subjects [mean: -1.87; $t(57)=-0.84$, $p=0.42$]. So — when analyzed separately — female, non-Hispanic and non-White subjects performed significantly better on paper-based tests than computer-based tests.

We then regressed the difference (C-P) onto the following predictors: age, OAT scores (academic average, total science and reading comprehension), undergraduate GPAs (total and math and science), GPAs in the first three semesters of our professional program, and learning styles. We found no significant correlations.

No validated questionnaire for determining attitudes toward paper and computerized examination formats is currently available. Therefore, we performed a factor analysis (via principal components) and found that the extracted communalities for all the survey question responses were significant (ranging from 0.65 to 0.96) and contributed to a single component that explained more than 37% of the variance in responses. It seems, therefore, that the survey questions represented, at least to some extent, student preference for one testing platform or the other. Therefore, we considered them all in our analysis.

The survey responses were not distributed normally, so we used a non-parametric test (Wilcoxon signed rank test) to determine whether the median responses were different than zero. A significant test and negative median indicated an overall preference (or endorsement) for the PBT format. If the median was positive, the endorsement was for the CBT. These results are summarized in **Table 3** and are shown for overall responses as well as broken down by gender and ethnicity (Hispanic or non-Hispanic). Subjects overall endorsed the PBT on 15 questions and the CBT on five questions. The results were identical for the 54 subjects who identified as non-Hispanic and similar for female subjects who endorsed paper examinations on two additional questions. Male subjects endorsed the computerized format for four questions but only endorsed paper for one question. Hispanic subjects endorsed paper testing for two questions and computer-based tests for one question.



Table 3. [Click to enlarge](#)

Some commonly endorsed items for paper were questions 1 (less stressed by paper exam), 5 (more positive experience with paper) and 22 (prefer to mark tentative answers and go back before recording final answer). Endorsements for computer-based testing included questions 4 (prefer the feedback after

taking the CBT), 8 (understand the need for CBT) and 18 (taking CBT prepares better for national licensure exams). To test whether these endorsements correlated in some way with a performance bias toward the paper or computerized formats, we created trichotomous variables for all survey questions, considering any negative response an endorsement for PBT, any positive response an endorsement for CBT, and any zero response to be “no opinion.” Because we are really only interested in whether these endorsements predicted significantly better performance on one testing platform or the other, we only analyzed subjects that scored at least 2.25 points higher on the computer-based test (those who performed best on the CBT) or at least 6.63 points better on paper (those who performed best on PBT; refer to Figure 4). We then calculated the likelihood (via X^2 analysis) that the distribution (number of PBT endorsements, no opinion, or CBT endorsements) was the same between these two performance groups.

There were significant findings for eight survey questions representing perceived difficulty (question 2), “mood” toward platforms (questions 9 and 17), overall preference (questions 6, 10, 13 and 15), and which platform better prepares for national licensure exams (question 18). Four representative significant findings are shown in **Figure 5**. To check for a “goodness of fit” (as in X^2 analysis), we sorted the scaled predictors into ordinal variables by assigning the lower quartile as “1”, second quartile as “2”, third quartile as “3”, and the upper quartile as “4.” Only the distributions of current optometry school GPA differed significantly between students performing best on PBT or CBT ($p=0.027$; **Figure 6**).



Figure 5. Each column represents the students who performed best (75th percentile) on the PBT (left; $n=14$) or the CBT (right; $n=14$). The different bands represent proportions whose attitudes favored CBT (black bands), PBT (light bands) or no preference (gray bands). The graphs shown are representative of significant findings regarding perceived difficulty (a), “mood” toward platforms (b), overall preference (c), and which platform better prepares them for national licensure exams (d). [Click to enlarge](#)



Figure 6. Each column represents the students who performed best (75th percentile) on the PBT (left; $n=14$) or the CBT (right; $n=14$). When examining all academic predictors, only distributions of current optometry school GPA differed significantly between these two groups. [Click to enlarge](#)

Discussion

This study found little difference in the performance of second-year optometry students on a computer-based test (CBT) compared with a paper-based-test (PBT) covering topics in organ pathology and ocular physiology. Overall, there was a trend toward higher mean scores on the PBT. In addition, an opinion survey found student attitudes favored the PBT more highly than the CBT.

Our examination performance findings are consistent with some but not all prior studies that have found little difference in student performance between CBT and PBT (see Bugbee¹⁹ and Vrabell²⁰ for reviews of the literature). Our results are consistent with those of Boevé et al.,²¹ who conducted a study with a crossover design somewhat like ours. In their undergraduate psychology course, one-half of the class ($n=199$) took their midterm exam on a computer while the other half of the class took the exam using paper and pencil. The groups were switched for the final examination. The researchers found no significant difference in the mean number of questions answered correctly between the computer-based and paper-based modes for both the midterm and final exam. In another recent study, Karay et al.²² evaluated the performance of 266 medical students on a standardized 200-question multiple-choice examination. Students were randomly assigned to take the exam on a computer or on paper. There was no significant difference in exam score between the groups, but students taking a PBT needed

significantly more time to complete the test.

Perhaps the trend towards PBT performance in our study was influenced by real or perceived difficulty differences between the courses or the individual examinations used in our design. There was a statistically significant difference between the mean scores for ocular physiology (84.5%) and organ pathology (81.1%). This difference could be explained by an enhanced interest in the ocular physiology topics among optometry students. However, the disparity between the mean PBT and CBT scores was larger (and statistically significant on paired sample t-tests) in the ocular physiology course (86.4% for PBT vs. 81.1% for the CBT; $p=0.003$) than in the organ pathology course where the PBT and CBT scores (81.5% and 80.6%, respectively) were essentially equivalent ($p=0.55$). It is quite possible that the PBT in the ocular physiology course covered material that was easier to master or had been introduced in a previous course. Regardless of the cause, our ability to draw inferences concerning any PBT bias may be weakened by the main effect of course and its interaction with exam mode on scores.

It has been suggested that males find technology more appealing and thus are more self-confident using computers.²³ In addition, there are indications that girls achieve less well than boys on computer-based problem-solving tasks.²⁴ This has led to a concern that females may perform worse on CBT than males. We did find that females performed significantly better on PBT than CBT, while for males there was no significant difference between exam modes. Our results agree with Jeong²⁵ who studied test scores of Korean grade-school children and found the CBT scores of female students to be significantly lower than their PBT scores in three of four subjects studied. This similarity should be considered in light of the notably different cohorts. Indeed, others have found no gender differences in more applicable investigations. For example, Clariana and Wallace²⁶ compared student performance on a 100-question multiple-choice examination. Fifty-four college students took the examination on a computer while 51 students took it on paper. No attempt was made to match the two groups for gender, academic achievement or any other variable, and the investigators found that gender was not significantly associated with computer vs. paper test mode effects.

Ethnic background and race are additional factors that may influence exam mode performance. It has been reported that grade-school children from an ethnic-minority background may have less exposure to computers both in the classroom and at home than children from the majority population.²³ While our study subjects overwhelmingly endorsed computerized testing in reporting they were "comfortable with technology," computer familiarity is one factor that has been identified as potentially influencing performance on CBT.²⁷ In our study, 10 students self-identified as being of Hispanic heritage and 54 who were non-Hispanic. Interestingly, we found that Hispanic students performed equally well on the CBT and PBT, while non-Hispanic subjects performed significantly better on the PBT. The 26 subjects who self-identified as non-White did perform significantly better on the PBT, but we found no performance bias among subjects who identified as White. These equivocal findings may indicate that disparities in computer familiarity may not be as much of a determining factor for this generation of students, at least not at the professional-school level.

No other individual student characteristic was significantly associated with exam mode performance in our analysis, in contrast with previous investigations. Watson²⁷ found that academically higher performing students benefited most from a computer-assisted learning program, while Clariana and Wallace²⁶ reported that higher attaining students performed significantly better on a CBT than a PBT, or conversely, PBT hindered the performance of high-attaining students more than low-attaining students. Our results may have been limited by a relatively low number of subjects; both previous studies involved nearly double the number subjects.

Following the second midterm examination, and after both midterm exam scores had been revealed to the students, each student completed a questionnaire. Responses from students with the greatest exam mode performance difference were analyzed for significant differences in their responses to survey

questions. In general, we found that students indicated a strong preference for the exam mode that they performed best on (Figure 5c). When asked which exam mode best prepares them for national licensure exams (which are CBT in optometry), nearly equal proportions of students acknowledged that CBTs were superior. This is not a surprising result, as NBEO Parts I and II (of III) are administered electronically. However, no student performing best on CBT acknowledged that PBT would better prepare them for licensing exams, and the overall distributions were significantly different (Figure 5d). These positive associations between performance and exam mode preference are not surprising. For example, it should be intuitive that more students who performed best on the paper-based test would think it was easier (Figure 5a) and that we should stay with that testing platform (Figure 5b). Our results differ from those of Washburn et al.¹³ who used a very similar design and found that students overwhelmingly preferred paper-pencil over computer-based assessments. However, the students in their study performed significantly better on the computer-based test.

We found a trend suggesting that differences in test anxiety may have contributed to differences in performance between the exam modes. While Washburn et al.¹³ found no association between test anxiety and test performance, it has been previously suggested that students with a lower comfort level with computers may experience greater test anxiety, and subsequent lower performance, with CBT.²⁸ In a study of 131 college undergraduate volunteers randomly assigned to computerized or paper-and-pencil versions of a battery of personality tests, Lankford et al.²⁹ found that female students and those with higher computer anxiety reported more depression when the test was administered on a computer. It is suggested that computer anxiety, like test anxiety in general, is not dependent on the degree of computer experience.²⁸ In our study, distribution differences in responses to questions such as "I was more stressed taking the test using..." and "I am more afraid of tests using..." approached significance, suggesting that anxiety levels were associated with exam mode performance difference. Perhaps our small study population and their foreknowledge of their exam performance prior to completing the questionnaire influenced our ability to detect a statistically significant correlation between anxiety and exam mode performance.

Strengths of our study include the 100% student participation rate, the crossover experimental design, and our access to student academic records, including undergraduate GPA and optometry school entrance exam performance. Furthermore, each student completed a VARK learning preference survey providing insight into their learning style.

Disadvantages that limit inferences from our study include the small sample size of only 64 students and a much smaller representation of Hispanic than non-Hispanic subjects. A further disadvantage is that to gain admission to optometry school students must perform well on the OAT, which is a CBT. Students that perform poorly on CBTs would not be expected to gain admission to optometry school. Although we found that OAT scores did not predict better performance on the CBT, we are dealing with a student population that is self-selected for good performance on CBT. One other disadvantage is that the two courses that were part of this experiment (organ pathology and ocular physiology) are not exactly equivalent. While the course material is similar (both covering foundational biologic principles) and were taught concurrently by the same instructor to the same students, there are differences in subject matter that may have influenced exam performance other than exam mode alone. For example, one course was 3 credit hours while the other was 2 credit hours. However, both courses were a continuation of first-year courses, and we believe that the difficulty of the course content was comparable throughout the semester within each course.

While some investigators have found no effect of question order on test performance,^{30,31} others have demonstrated effects on scores³² and score distributions.³³ In the current study, PBTs were administered in one version with a fixed order of questions and answer choices. There were, however, multiple versions of CBTs, which limits our ability to draw inferences about differences between the two formats. Lastly, we need to acknowledge that a better investigational approach may have been to administer

each of the four midterms in both formats: PBT and CBT. Students could have been randomized into CBT and PBT for each exam and swapped for the second in each course. We decided against this approach for instructional reasons as students may have considered it unfair to take their first CBT while others in their class were being evaluated over the same material with the more familiar PBT.

Conclusions

In summary, we found no statistically significant difference in overall performance between PBT and CBT in this group of healthcare professional students. Females, non-White students, and non-Hispanic students performed significantly better on PBT. We also found that performance differences between the two formats predicted student perceptions of difficulty, preference and utility of the formats. In addition, we found trends suggesting that test anxiety may contribute to poor CBT performance among some students. These trends warrant further investigation. We plan to conduct a follow-up study on this same cohort of students to examine how their attitudes and beliefs toward CBT may have evolved with increased familiarity with this mode of exam administration.

Acknowledgement

This research was supported by an Innovation in Teaching grant from the American Academy of Optometry Foundation and Johnson & Johnson Vision.

References

1. Shepard LA. The role of assessment in a learning culture. *Educational Researcher*. 2000;29(7):4-14.
2. Association of Schools and Colleges of Optometry (ASCO). Attributes of Students Graduating from Schools and Colleges of Optometry: A 2011 Report from the Association of Schools and Colleges of Optometry. 2011 [cited Aug 26, 2019]; Available from: <https://optometriceducation.org/faculty-and-administrators/asco-policy-guidelines-and-reports/>.
3. Accreditation Council on Optometric Education (ACOE). Professional Optometric Degree Standards. 2016 [cited Aug 26, 2019]; Available from: <https://www.aoa.org/optometrists/for-educators/accreditation-council-on-optometric-education/accreditation-resources-and-guidance/optometric-degree-programs->
4. National Board of Examiners in Optometry (NBEO). Exam Content Outlines. 2019 [cited Aug 26, 2019]; Available from: https://www.optometry.org/exam_content.cfm
5. Shavelson RJ. Assessing student learning responsibly: from history to an audacious proposal. *Change*. 2007; 39(1):26-33.
6. Pellegrino JW. The evolution of educational assessment: considering the past and imagining the future. The sixth annual William H. Angoff Memorial Lecture. Presented at: Educational Testing Service; Nov 17, 1999; Princeton, New Jersey. 2004.
7. Kuikka M, Kitola M, Laakso M. Challenges when introducing electronic exam. *Res Learn Technol*. 2014;22:22817 (e-copy). doi: 10.3402/rlt.v22.22817
8. Bussieres J-F, Metras M-E, Leclerc G. Use of Moodle, ExamSoft, and Twitter in a first-year pharmacy course. *Am J Pharm Educ*. 2012;76(5):94.
9. Cook J, Jenkins V. Getting started with e-assessment. Bath: University of Bath. 2010.
10. Craven P. History and challenges of e-assessment: The 'Cambridge Approach' perspective – e-assessment research and development 1989-2009. Cambridge: Cambridge University. 2009.
11. Millsap CM. Comparison of computer testing versus traditional paper-and-pencil testing [dissertation]. Denton (TX): University of North Texas. 2000.
12. Capay M, Magdin M, Mesarosava M. Enhancement of e-testing possibilities with the elements of interactivity reflecting the students' attitude to electronic testing. Paper presented at: Proceedings of the 10th European Conference on e-Learning; Nov 4-5; Brighton UK. 2011.

13. Washburn S, Herman J, Stewart R. Evaluation of performance and perceptions of electronic vs. paper multiple-choice exams. *Adv Physiol Educ.* 2017;41:548-555.
14. Lovelace M, Brickman P. 2013. Best practices for measuring students' attitudes toward learning science. *CBE Life Sci Educ.* 2013 Winter;12(4):606-617.
15. Akhlaghi N, Mirkazemi H, Jafarzade M, Akhlaghi N. Does learning style preferences influence academic performance among dental students in Isfahan, Iran? *J Educ Eval Health Prof.* 2018 Mar 24;15:8. doi:10.3352/jeehp.2018.15.8
16. American Psychological Association. Guidelines for computer-based tests and interpretations. Washington, DC; 1986.
17. Bland J, Altman D. Measuring agreement in method comparison studies. *Stat Methods Med Res.* 1999;8(2):135-160.
18. Bland J, Altman D. 1986. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet.* 1986;327(8476):307-310.
19. Bugbee AC. The equivalence of paper-and-pencil and computer-based testing. *Journal of Research on Computing in Education.* 1996;28(3):282-299.
20. Vrabel M. Computerized versus paper-and-pencil testing methods for a nursing certification examination: a review of the literature. *Comput Inform Nurs.* 2004;22(2):94-98.
21. Boevé AJ, Meijer RR, Albers CJ, Beetsma Y, Bosker RJ. Introducing computer-based testing in high-stakes exams in higher education: results of a field experiment. *PLoS ONE.* 2015;10(12):e0143616. doi: 10.1371/journal.pone.0143616.
22. Karay Y, Schaubert SK, Stosch C, Schuttpelz-Brauns K. Computer versus paper—does it make any difference in test performance? *Teach Learn Med.* 2015;27(1):57-62.
23. Volman M, van Eck E, Heemskerk I, Kuiper E. New technologies, new differences. Gender and ethnic differences in pupils' use of ICT in primary and secondary education. *Computers & Education.* 2005;45(1):35-55.
24. Barbieri MS, Light PH. Interaction, gender, and performance on a computer-based problem solving task. *Learn Instr.* 1992;2(3):199-213.
25. Jeong H. A comparative study of scores on computer-based tests and paper-based tests. *Beh Inform Technol.* 2014;33(4):410-422.
26. Clariana R, Wallace P. Paper-based versus computer-based assessment: key factors associated with the test mode effect. *Brit J Educ Technol.* 2002;33(5):593-602.
27. Watson B. Key factors affecting conceptual gains from CAL materials. *Brit J Educ Technol.* 2001;32:587-593.
28. Marcoulides GA. The relationship between computer anxiety and computer achievement. *Educ Comput Res.* 1988;4(2):177-187.
29. Lankford JS, Bell RW, Elias JW. Computerized versus standard personality measures: equivalency, computer anxiety, and gender differences. *Computers in Human Behavior.* 1994;10(4):497-510.
30. Monk JJ, Stallings WM. Effects of item order on test scores. *J Ed Research.* 1970;63(10):463-465.
31. Gohmann SF, Spector LC. Test scrambling and student performance. *J Econ Ed.* 1989;20(3):235-238.
32. Gruber RA. Sequencing exam questions relative to topic presentation. *J Acct Ed.* 1987;5(1):77-86.
33. Carlson JL, Ostrosky AL. Item sequence and student performance on multiple-choice exams: further evidence. *J Econ Ed.* 1992;23(3):232-235.

Brian K. Foutch, OD, PhD, [foutch@uiwtx.edu] is an Associate Professor at the University of the Incarnate Word (UIW) Rosenberg School of Optometry where he teaches epidemiology, public health and neuroanatomy. He also precepts third- and fourth-year optometry interns in Primary Care clinics at the UIW Eye Institute.

Lourdes Fortepiani, MD, PhD, is an Associate Professor at the University of the Incarnate Word

Rosenberg School of Optometry where she teaches foundational sciences including physiology, systemic disease, ocular physiology and ocular correlates of systemic disease. She is also an Adjunct Assistant Professor in the Department of Cellular and Integrative Physiology at UT Health San Antonio.

Richard Trevino, OD, is an Associate Professor at the University of the Incarnate Word (UIW) Rosenberg School of Optometry where he teaches advanced clinical procedures, glaucoma and posterior segment diagnosis and management. He also serves as Chief of the Ocular Disease Service at the UIW Eye Institute.

PEER REVIEWED

Development of a Polymeric Eye Model for Foreign Body Removal

Chau-Minh Phan, PhD, MSc, FAAO, Hendrik Walther, PhD, MSc, FAAO, and Lyndon Jones, PhD, DSc, FCOptom, FAAO | Optometric Education: Volume 45, Number 1 (Fall 2019)

Abstract

Background: Removal of a corneal foreign body is a critical skill for optometrists and ophthalmologists to practice, but most students receive minimal and rudimentary training in this area. The aim of this study was to develop a simple polymeric eye model, with embedded metal particles, for teaching foreign body removal. **Methods:** The eye models were made from 15% weight/volume polyvinyl alcohol, molded into an eyeball shape and embedded with 100- 300- μm size steel particles. The model was tested with 90 third-year optometry students, and a follow-up survey was conducted to rate the usefulness and realism of the model. **Results:** Of the 33 survey respondents, the majority (70%) thought the model was both useful and realistic. **Conclusion:** Students can easily use this model to improve their hand-eye coordination and confidence in removing corneal foreign bodies and rust rings under a slit lamp microscope.

Key Words: foreign body removal, eye model, polyvinyl alcohol, corneal foreign body removal, FBR

Introduction

Corneal foreign bodies, such as particulates of metal, glass, wood, plastic or sand, are among the most common causes of ocular injury.¹⁻³ When left untreated, they can potentially lead to tissue death, infections and vision loss.^{1,2} Unfortunately, physicians are often poorly trained in the removal of corneal foreign bodies, which may lead to delayed or suboptimal treatment in the emergency room setting.³

The best approach to removing foreign bodies is under slit lamp magnification using a sterile large-gauge needle, followed by removal of the rust ring with an electric burr (AlgerBrush).^{1,2} However, the slit lamp's degree of magnification and changes in depth of field can lead to exaggerated hand movements. The removal technique therefore requires precise hand-eye coordination and extremely fine motor skills, both of which are acquired through extensive practice.^{4,5} For this reason, adequate instructions and training for corneal foreign body removal (FBR) is critical in ophthalmic and optometric programs.^{3,6,7} Regrettably, there are not enough opportunities for students to gain the necessary practical experience.³

Studies have shown that students practicing FBR on simulated eye models improve their confidence levels and skills in that particular art.³⁻⁵ One of the first simulated models, and perhaps the most common, are bovine eyes with embedded metal pieces from an angle grinder.³ The embedded metal forms rust rings closely resembling those found in the human cornea. This provides a fairly realistic simulation of foreign bodies.³ The bovine eyes can also be substituted for porcine eyes, depending on the availability from local slaughterhouses.

Deceased animal eyes are relatively cost-effective and readily available. Furthermore, the anatomies of porcine eyes and bovine eyes are relatively similar to that of human eyes. However, they are biological tissue. As such, disposal of these eyes, and the post-training disinfection process for the tools used, must follow rigorous procedures for biological hazards. The dead tissue will also release unpleasant

odors. As a result, an alternative non-animal eye model is preferred.

Several approaches to using non-animal eye models for FBR have been tried. In general, these methods use a readily available polymer, such as paraffin wax, agar or ballistic gel, which is molded into the shape of an eye.^{4,5,8} The foreign bodies, consisting of various small pieces, are embedded afterwards. However, these polymers do not mimic the consistency of the cornea, which has a high water content as well as a mechanical strength. Studies have shown that polyvinyl alcohol (PVA) can be used to create an artificial human cornea and vitreous humor.^{9,10} PVA is ideal due to its low cost, ready availability, high water content (close to that of the cornea) and excellent mechanical properties.¹¹⁻¹⁴ It is commonly used in medical devices due to its favorable biocompatibility, chemical resistance and low protein adsorption properties.¹⁴ The advantages of using a PVA eye model for FBR are that students can safely use their own equipment to practice the procedure, there are no rigorous disinfection procedures, and the “eyes” can be kept in storage for long periods of time without spoilage. To our knowledge, eye models using PVA have not been studied. The aim of this study was to create a simple eye model, using PVA, with embedded metal particles that would assist in the teaching of FBR.

Methods

Molds

The molds for the eyeballs were designed using a computer-aided design (CAD) program and were computer numerical control (CNC) machined from acetal plastic. The dimensions for the eye were 22.5 mm x 22.5 mm x 24 mm for height x width x axial length. The molds consisted of two parts, the anterior half for the front half of the eye, and the posterior half for the back of the eye (**Figure 1**). Additionally, the posterior mold had two holes to allow for filling and the escape of any excess polymer solution. A quarter-inch diameter acrylic rod was used to create a cylindrical hole inside the eyeball to allow for mounting during FBR training.



Figure 1. A: Custom computer numerical control (CNC)-machined molds for the anterior half of the eyeball, the posterior half of the eyeball, and a quarter-inch acrylic rod used to make a hole in the eyeball. **B:** The molds assembled.

Synthesis of polyvinyl eyeball

Dimethyl sulfoxide, and PVA with a molecular weight of (M_w) 89-98 kilodaltons (kDa) (99%+ hydrolyzed), were purchased from Sigma Aldrich (St. Louis, Mo.). The procedure was adopted from methods proposed by Hyon et al.¹² In brief, PVA was added to a mixture of dimethyl sulfoxide and Milli-Q water (8:2) to achieve a concentration of 15% weight/volume (w/v). Milli-Q water, commonly available in most laboratories, is ultrapure water obtained from filtering distilled water through a Millipore ion exchange system. The concentration can be increased if a harder eyeball is desired. The mixture was then lightly stirred for 5 minutes, and then heated in an oven at 115-120°C for 3 hours. After the heating step, the PVA turned clear and viscous, at which point it was poured into the eye mold and the acrylic rod was pushed into the back of the posterior mold to form a cylindrical mounting hole. Subsequently, the mold was held at -30°C for 3 hours for gelation to occur. The gelled eyeballs were then removed from the molds and placed in 500 mL of Milli-Q water, renewed daily, for 3 days to remove the organic solvents.

Foreign bodies

A universal milling machine was used to obtain fine steel particulates from a block of steel metal. The steel particulates were then filtered using 100 μ m x 100 μ m, and 300 μ m x 300 μ m, nylon Spectra Mesh Woven Filters to obtain particle sizes between 100 μ m and 300 μ m. Several fine particulates of steel were placed in the front half of the eyeball molds and heated at 115-120°C for 15 minutes. The molds

were then removed from the oven, and the PVA eyeballs were pressed against the front half of the mold so the front surface of the polymer eyeball re-melted and the heated steel particulates could melt onto the front surface of the eye. The eyeball and the molds were frozen immediately at -30°C for 30 min for re-gelation. The eyeballs were then removed from the molds and stored in Milli-Q water until use. After approximately 24 hours, the steel particulate naturally formed a rust ring in the model eye.

Mount structure

A custom mount to secure the eyeball to a slit lamp headrest was designed using a CAD program. The mount was then cut from quarter-inch acrylic sheets, sourced from the local engineering workshop, using a laser cutter. The structures were assembled and glued together using methylene chloride. A quarter-inch acrylic rod was cut in a 2-mm length to easily mount and dismount the eyeball to the structure.

Students

A total of 50 eye models were used and tested by 90 third-year optometry students during their FBR lab. After the lab, an online survey was sent to all students asking them to rate the usefulness and realism of the simulated eye model for FBR training.

Results

The model eyeball with embedded metal foreign bodies and rust rings are shown in **Figure 2**. The magnified view under the slit lamp clearly shows the foreign bodies and rust rings. The PVA at 15% w/v concentration has a semi jelly-like consistency that is also resistant to puncturing. **Figure 3** shows the custom acrylic structure used to mount the eyeball to a standard slit lamp headrest. Students can easily practice corneal FBR and rust ring removal using the eye model, as shown in **Figure 4**.

Of the 90 third-year optometry students who received the online survey after the FBR lab, 33 responded. More than 72% of the respondents thought the eye models were either very useful (48.48%) or extremely useful (24.24%) in learning FBR, while 70% of the respondents thought the models were fairly realistic (**Figure 5**).



Figure 2. A: Polyvinyl alcohol eyeball model with embedded foreign bodies and rust ring. **B:** The eyeball model viewed through a slit lamp under magnification.

[Click to enlarge](#)



Figure 3. Custom acrylic structure for mounting eyeball to a slit lamp.

[Click to enlarge](#)



Figure 4. A: A student practicing corneal foreign body removal. **B:** A student practicing rust ring removal.

Discussion

The present study provides a simple method for creating eye models containing corneal foreign bodies. Multiple iterations of this eye model were vetted, tested and approved by the faculty members in charge of administrating the FBR lab. The final version of the eye model was tested on 90 third-year optometry students, and based on the survey, the overall response to this model was enthusiastic. More than 72% of the students thought the model was very useful, and 70% of the students thought the model was fairly realistic. The students initially found it challenging to remove the foreign bodies while under magnification from the slit lamp, but after a few attempts they quickly gained confidence in the FBR

procedure.



Figure 5. Survey responses from 33 optometry students regarding the realism **(A)** and usefulness **(B)** of the eye model for learning foreign body removal.

[Click to enlarge](#)

The most common conventional eye models for teaching foreign body removal are bovine or porcine eyes with embedded metal filings.³ In the past, we had used porcine eyes, but the disinfection process post-procedure for dead animal eyes was time consuming and rigorous, and the foul odors were undesirable. We also found that it was difficult to consistently create large enough foreign bodies that were also isolated enough from each other using an angle grinder.³ Often there would be numerous clusters of extremely fine particles, which could be easily rinsed off with water.

Considering the importance of corneal FBR, very few studies have been published on alternative models to animal eyes.^{4,5,8,15} Austin et al. created an eye model using marbles dipped in melted paraffin. The foreign bodies were made from snipping small pieces from an 18-gauge copper wire, and the rust rings were simulated using a rust-colored crayon.⁴ Another model, proposed by Cheng et al., used expired agar plates, readily available from the local microbiology departments.⁵ Materials such as gravel were used as the simulated foreign bodies.⁵ In the past, we have tried both paraffin wax and agar, but found that both materials did not mimic the consistency of corneal tissue. Paraffin wax was too hydrophobic and sticky, while agar was too brittle and soft, and consequently the foreign bodies were removed too easily. An interesting model was proposed by Sayegh et al. It used ballistic gel made from silicone and corn starch.⁸ Ballistic gels are a class of gels that mimic the consistency of muscle tissue, and they are typically used to simulate bullet wounds.¹⁶ The foreign bodies were made from paper clips snipped in 2-mm portions using a wire cutter.⁸

We considered two important factors for creating a realistic eye model for FBR. The first was the consistency of the eyeball, which needed to have a high water content and resistance similar to an actual cornea. We chose PVA because of its low cost, high water content¹³ and excellent tensile strength.¹¹ Studies also have shown that it can be used as a material for artificial corneas and human vitreous humor.^{9,10} The material synthesis process that we proposed is also flexible. For instance, the concentration of PVA can be increased or decreased to obtain a stronger or weaker gel. Additionally, this approach to synthesizing PVA gels is thermo-reversible, which allows melting of the surface of the gels to incorporate the foreign bodies.

The second important factor we considered was the quality of the foreign body, which needed to be similar to the size of a foreign body typically seen in practice. In addition, the foreign body had to be embedded only on the surface, yet deep enough into the gel to be comparable with what is seen in real-life situations. To ensure similar sizes of our foreign bodies, we obtained filings of various dimensions from a steel block using a universal milling machine and then filtered the filings through 100- μ m- and 300- μ m nylon filters to obtain particle sizes between 100 μ m and 300 μ m. We embedded the particles by heating the metal filings to 120°C before pressing/melting them into the eyeball. By freezing the eyeballs rapidly afterwards, the foreign bodies became tightly embedded within the upper layers of the PVA material.

One of the main drawbacks of our eye model for FBR is the lack of Bowman's membrane to resist the penetration of the needle. Without this layer, a student is able to pierce deeper into the eye model than would normally be possible in a real eye. This layer could be simulated by incorporating another thicker layer, made from a higher concentration of PVA or a different polymer such as polydimethylsiloxane.

Future studies will examine the synthesis of multiple layers representing different structures of the eye for a more realistic simulation.

Conclusion

This study tested a simple and versatile method for building an eye model to aid in teaching corneal foreign body removal. The eye model is made from polyvinyl alcohol molded into the shape of an eye and embedded with steel particles. The overall eye model has very high wettability and similar consistency to an actual eye, and the foreign bodies also form a natural rust ring. Students can easily use this model with a slit lamp to learn the necessary techniques to remove foreign bodies and rust rings. The advantages of using this model over the traditional bovine or porcine eye is a simpler cleanup, students can use their own equipment to practice, the eyes do not spoil in storage, and there is no unpleasant odor.

References

1. Luo Z, Gardiner M. The incidence of intraocular foreign bodies and other intraocular findings in patients with corneal metal foreign bodies. *Ophthalmology*. 2010;117(11):2218-21.
2. Ramakrishnan T, Constantinou M, Jhanji V, Vajpayee RB. Corneal metallic foreign body injuries due to suboptimal ocular protection. *Arch Environ Occup Health*. 2012;67(1):48-50.
3. Collins DW, Coroneo MT. Removal of corneal foreign bodies: an instructional model. *Ophthalmic Surg*. 1994;25(2):99-101.
4. Austin PE, Ljung M, Dunn KA. A new model for teaching corneal foreign Body removal. *Acad Emerg Med*. 1995;2(9):831-4.
5. Cheng ML, Fu L, Cackett P. A novel, safe and cost effective way for teaching corneal foreign body removal. *Emerg Med J*. 2015;32(6):501-2.
6. Spivey BE. Ophthalmology for medical students: content and comment. *Arch Ophthalmol*. 1970 Sep;84(3):368-75.
7. McCarthy W, Donaldson E, Hollows F. What should we teach? A consensus method to determine curriculum content. *Medical Education*. 1977 Jul;11(4):249-56.
8. Sayegh JS, Lahham S, Woodhouse L, Seong J, McCoy CE. Creation of a realistic model for removal of a metallic corneal foreign body for less than \$75. *West J Emerg Med*. 2017;18(1):121-25.
9. Leone G, Consumi M, Aggravi M, Donati A, Lamponi S, Magnani A. PVA/STMP based hydrogels as potential substitutes of human vitreous. *J Mater Sci Mater Med*. 2010;21(8):2491-500.
10. Wang J, Gao C, Zhang Y, Wan Y. Preparation and in vitro characterization of BC/PVA hydrogel composite for its potential use as artificial cornea biomaterial. *Mater Sci Eng C Mater Biol Appl*. 2010;30(1):214-8.
11. Stammen JA, Williams S, Ku DN, Guldberg RE. Mechanical properties of a novel PVA hydrogel in shear and unconfined compression. *Biomaterials*. 2001 Apr;22(8):799-806.
12. Hyon SH, Cha WI, Ikada Y, Kita M, Ogura Y, Honda Y. Poly (vinyl alcohol) hydrogels as soft contact lens material. *J Biomater Sci, Polym Ed*. 1994;5(5):397-406.
13. Hyon SH, Cha WI, Ikada Y. Preparation of transparent poly(vinyl alcohol) hydrogel. *Polym Bull*. 1989;22(2):119-22.
14. Baker MI, Walsh SP, Schwartz Z, Boyan BD. A review of polyvinyl alcohol and its uses in cartilage and orthopedic applications. *J Biomed Mater Res B Appl Biomater*. 2012 Jul;100(5):1451-7.
15. Radow BK. Variable pathological and surgical eye model and method related thereto. United States patent US5893719A. 1999 Apr 13.
16. Jussila J. Preparing ballistic gelatine – review and proposal for a standard method. *Forensic Sci Int*. 2004 May 10;141(2-3):91-8.

Dr. Phan [chau.minh.phan@gmail.com] is a post-doctoral fellow at the Centre for Ocular Research & Education (CORE), School of Optometry and Vision Science, at the University of Waterloo in Waterloo, Ontario, Canada. He has a keen interest in entrepreneurship and commercialization, having launched two start-ups, Tricolops Technology and OcuBlink Inc. His research interests primarily focus on development of novel ophthalmic materials and physiological eye models.

Dr. Walther began his career in optometry as an optician in Germany, undertaking an apprenticeship (2002-2005). He later received BSc (2009) and MSc (2011) degrees in optometry from the University of Applied Sciences in Jena. Afterwards, he joined the University of Waterloo to complete his doctorate (2017), investigating ways to quantify lipid deposition onto various contact lens materials. He is currently a post-doctoral fellow at the Centre for Ocular Research & Education (CORE), School of Optometry and Vision Science, at the University of Waterloo in Waterloo, Ontario, Canada.

Dr. Jones [lwjones@uwaterloo.ca] is a Professor at the School of Optometry and Vision Science, University Research Chair and Director of the Centre for Ocular Research & Education (CORE) at the University of Waterloo in Waterloo, Ontario, Canada. His research interests primarily focus on the interaction of novel and existing contact lens materials with the ocular environment, dry eye and the development of novel materials for ocular drug delivery.

PEER REVIEWED

Lymphoproliferative Disorders of the Ocular Adnexa: a Teaching Case Report of Conjunctival MALT Lymphoma and Lymphoid Hyperplasia

Ellen McCrary, OD, Andrea Yiasemis, OD, FAAO, and Pauline F. Ilsen, OD, FAAO | Optometric Education: Volume 45, Number 1 (Fall 2019)

Abstract

Ocular adnexal lymphoproliferative disease (OALD) refers to a spectrum of lymphoid disorders affecting the eyelid, conjunctiva, lacrimal gland and orbit. They may be indistinguishable on clinical examination and imaging. Thus, further evaluation is necessary to characterize the lesions and determine whether they are localized or systemic. To highlight the spectrum of disorders, two cases of OALD are presented: one case of benign lymphoid hyperplasia and one case of malignant lymphoma. Diagnosis, management and prognosis of these conditions are discussed. Eyecare providers should be aware of the clinical presentation, diagnostic testing and treatment of the lymphoproliferative disorders as they may present in patients who have no known history of lymphoma and may be readily mistaken for other conditions.

Key Words: lymphoproliferative disorder, lymphoma, benign reactive lymphoid hyperplasia, MALT lymphoma

Background

Ocular adnexal lymphoproliferative disorders (OALD) range from the benign reactive lymphoid hyperplasias (RLHs) to the malignant lymphomas. Both benign and malignant ocular lymphoproliferative disorders typically affect the conjunctiva, orbit and lacrimal gland and are most often unilateral.¹⁻⁵ OALD has a slight female predilection and typically presents between the 5th and 7th decades.^{4,6} Malignant lymphomas and benign lymphoid hyperplasias (LHs) often appear identical on clinical examination and imaging.

We present two cases of OALD that initially appeared similar on clinical examination, but after further evaluation with biopsy, immunohistochemistry and molecular genetics were found to be respectively malignant and benign. Local and whole-body imaging is needed to determine whether lymphoid disease is localized to the orbit or secondary to systemic disease.

This teaching case report is aimed at fourth-year optometry students, residents and current practitioners. Eyecare providers should be aware of the clinical presentation, diagnostic techniques, management options and outcomes for the lymphoproliferative disorders of the ocular adnexa as they may present without prior known history of lymphoma. OALD may be easily mistaken for other conditions, and the eyecare provider may play a significant role in co-management with other disciplines.

Case Descriptions

Case report 1: MALT lymphoma of the palpebral conjunctiva

A 61-year-old black male presented for a routine eye examination with a chief complaint of blurry vision at distance and near. His last eye exam was 35 years ago and the only vision correction he was wearing

were over-the-counter readers. He had no history of eye injury or surgery and no personal or family history of ocular disease. His medical history was positive for hypertension, a benign liver mass, hepatitis C, chronic obstructive pulmonary disease, peptic ulcer disease, anemia and past substance abuse. His best-corrected visual acuity was 20/15 for both the right and left eye.

Slit-lamp biomicroscopy revealed a large, red, fleshy and gelatinous conjunctival mass in the left eye that was present in almost the entire inferior fornix, palpebral conjunctiva and encroaching onto the bulbar conjunctiva. The patient was unaware of the mass and was asymptomatic. Lymphoproliferative disease was suspected, and photographs were taken at the initial visit (**Figure 1A**). All other examination findings were unremarkable.

The patient was referred to the ophthalmology department for evaluation but did not follow up until the next year. At time of presentation with ophthalmology one year later, he was still mostly asymptomatic, although he reported an occasional mild foreign body sensation. The left lower lid now exhibited a slight fullness, and his blink was abnormal with a tendency toward entropion. The pink conjunctival mass involving most of the lower lid and fornix measured approximately 12 mm x 6 mm on the bulbar conjunctiva and 25 mm horizontally on the palpebral conjunctiva. A complete blood count (CBC) with differential was ordered, and the patient was referred to the oculoplastics clinic for biopsy. The CBC revealed a mild elevation of monocytes and basophils but was stable compared with baseline testing over the past 14 years.



Figure 1. A 61-year-old black male with MALT lymphoma of the inferior palpebral conjunctiva and fornix before **(A)** and 2 years after **(B)** treatment with radiotherapy. [Click to enlarge](#)

An incisional biopsy was performed one month later. Histology revealed “a uniform population of lymphocytes with occasional lympho-epithelial islands within the subepithelial tissue.” Flow cytometry showed CD19+, CD20+ and CD22+ cell markers and lambda light chain restriction. A diagnosis of B-cell lymphoma consistent with mucosa-associated lymphoid tissue (MALT) lymphoma was made. Positron emission tomography (PET) showed no systemic involvement, and the final diagnosis was stage IE extranodal B-cell marginal-zone lymphoma of the left inferior conjunctiva. Intensity-modulated radiation therapy (IMRT) was performed with a total of 36 Gy split into 18 fractions over 26 days. The conjunctival mass completely resolved without recurrence at the 9-month follow-up exam.

The patient returned to clinic 2 years later with a complaint of worsening vision in the left eye and glare with nighttime driving. Vision was correctable to 20/20 in the right and left eyes but with glare testing declined to 20/80 in the left eye. Asymmetric cataracts were found, with a denser, milky nuclear sclerotic cataract and a small posterior subcapsular cataract in the left eye. There was no evidence of MALT lymphoma recurrence or systemic involvement (**Figure 1B**).

Case report 2: lymphoid hyperplasia of the palpebral conjunctiva and fornix

A 68-year-old white male known to the eye clinic presented with a complaint of a “bump” inside his left lower lid for the past 10 days. He noted that initially the lesion felt hard and small, but 4 days prior to the exam it had become flatter and more spread out. The patient had no pain or other associated symptoms. His medical history was positive for hyperlipidemia, hearing loss, erectile dysfunction, chronic kidney disease, migraines and basal cell carcinoma on the left cheek that was removed 1.5 years prior. His last eye exam was 6 months ago at the same clinic and was unremarkable. Best-corrected visual acuity was 20/20 in the right eye and 20/25+2 in the left eye.

Slit-lamp examination revealed a 12-mm horizontal by 5-mm vertical firm, fleshy mass encompassing

almost the entire lower palpebral conjunctiva and fornix with no tenderness on palpation. The remainder of the examination was unremarkable. An atypical chalazion was suspected and aggressive hot compresses were initiated. A 1-week follow-up visit was scheduled due to the sudden onset and atypical fleshy appearance of the lesion.



Figure 2. A 68-year-old white male with benign reactive lymphoid hyperplasia of the inferior palpebral conjunctiva and fornix. [Click to enlarge](#)

At the 1-week follow-up, the patient reported that the lesion was reduced in size, but examination revealed spreading of the lesion more nasally and into the fornix and inferior bulbar conjunctiva. Best-corrected visual acuity was stable in both eyes. Photographs of the lesion were taken, and the patient was referred to the oculoplastics clinic on the same day for evaluation (**Figure 2**). The lesion was biopsied, and a pathology evaluation was performed. Immunohistochemistry showed “mixed CD3 and CD20 lymphocytic populations with negative bc12 stain on germinal centers and positive CD10 stain almost exclusively by the follicular center cells.” Immunophenotyping showed a mixture of 36% T cells (CD3+) and 62% B cells (CD19+ and CD20+) and an elevated CD4:CD8 ratio of 6:1. A diagnosis of benign RLH was made.

When the patient returned to the oculoplastics clinic 2 weeks later, the treatment options were discussed with him and he chose observation only at that time. A plan was made to monitor the lesion every 3 months and to consider steroid injections in the future if any growth occurred. However, the patient was lost to follow-up for 8 months. When he returned, the conjunctival lesion was not appreciated. The patient reported that the lesion had resolved on its own approximately 3 weeks after being biopsied.

Education Guidelines

Learning objectives

1. Understand the spectrum of ocular adnexal lymphoproliferative disease and be familiar with its typical appearance
2. Be aware of differential diagnoses for lymphoproliferative disorders
3. Become familiar with how to manage a patient with suspected ocular adnexal lymphoproliferative disease
4. Become familiar with traditional and emerging immunotherapy treatments for ocular adnexal lymphoproliferative disease

Key concepts

1. Recognize that RLH is clinically indistinguishable from ocular lymphoma
2. Recognize that lymphoproliferative disease is often asymptomatic; however, clinicians should be familiar with common signs and symptoms
3. Become familiar with appropriate workup of suspected ocular lymphomas, including imaging and histology

Discussion points

1. What is the clinical presentation of OALD? What are the differential diagnoses?
2. What additional tests/workup are indicated when OALD is suspected?
3. What is the pathogenesis of lymphoproliferative disease?
4. What is the treatment for OALD?
5. What is the prognosis for OALD?

Literature review

Ocular adnexal lymphoproliferative disease refers to a spectrum of lymphoid disorders affecting the eyelid, conjunctiva, lacrimal gland and orbit. Histopathology classifies the ocular adnexal lymphomas as either benign RLH or malignant lymphoma.⁷

Ocular adnexal lymphoma (OAL) is the most common type of OALD and accounts for 68-98% of all lymphoid lesions.^{1,6,8} OAL is considered primary if it is located solely in the ocular tissues, while secondary lymphoma represents metastasis from another location in the body that is found either subsequently or simultaneously to the systemic lesions.^{2,9-10} OAL is further categorized into Hodgkin disease and non-Hodgkin lymphoma (NHL).

The majority of OAL is non-Hodgkin B-cell lymphoma, the most common of which is extranodal marginal-zone lymphoma, commonly referred to as MALT lymphoma (46-75%). This type of lymphoma was highlighted in the first case description. Other common types of OAL include follicular lymphoma (4.5-18%), diffuse large-B-cell lymphoma (4.7-16%) and mantle-cell lymphoma (3-9%).^{2-3,11-13} Studies have either found no gender predilection, or a slight female majority.^{2-4,13-16} The lymphoproliferative disorders are most common in older adults, with a median age of 59-71.^{1,4,8,13} Over the past few decades, OALD has been on the rise in Western countries. Studies show annual increases of 3.4-6.5%, mainly due to an increase in ocular adnexal MALT lymphoma.^{13,16} This increase may be due to advances in diagnostic techniques but does not account for the fact that the increase of systemic NHL has already peaked while OAL is still increasing.¹³

RLH is a proliferation of lymphoid tissue, usually with a polyclonal mix of small lymphocytes, and is a benign condition.¹⁴ It has a predilection for the orbit and adnexa but only accounts for approximately 7-15% of OALD.^{1,8,11-12,15} RLH is essentially indistinguishable from NHL clinically, and it requires biopsy for diagnosis as demonstrated in the second case description.

Discussion

Teaching instructions: Participants should read each question and consider how they would respond and then read the information provided in the text. Learning objectives are to be assessed by comparing participants' responses to the information provided.

What is the clinical presentation of ocular adnexal lymphoproliferative disease? What are some differential diagnoses?

Lymphoma is most common in the 5th to 7th decades.^{4,6} LH typically presents 5-10 years earlier.^{3,6} Both of the patients in the cases presented were in their 6th decade. Lymphoproliferative disease has a slight female predominance.^{1,3-5} The clinical appearances of benign LH and malignant lymphoma are also similar, and they are indistinguishable on clinical examination.^{1-2,6} Lymphoproliferative disease can be asymptomatic and found on routine exam but often presents as a slowly enlarging, painless mass. Common symptoms include proptosis, swelling, diplopia, ptosis, and mild to no pain and inflammation.^{1-2,5,11,17} Vision loss is rare as these lesions mold to the globe and orbit rather than invade the surrounding tissues.¹⁷⁻¹⁸ The median duration of symptoms before diagnosis is 6-7 months but can range from 1 month to 10 years.^{4,11}

The most common locations involved in lymphoproliferative disorders of the ocular adnexa are the orbit/lacrimal gland (33-46%), conjunctiva (23-42%) and eyelid (10-25%).¹⁻⁴ Conjunctival location usually offers the best prognosis, while eyelid lymphoma is usually secondary to a more aggressive systemic lymphoma,^{3,15} and lacrimal lymphoma is most likely to spread to the lymph nodes.¹⁹ The majority of OALD is unilateral; only 10-25% is bilateral, which often signifies a poorer prognosis.^{4,12,17}

Orbital and lacrimal gland lesions usually present as a firm or rubbery mass that can lead to proptosis.⁴ Conjunctival lymphoproliferation is often a characteristic “salmon-colored” lesion that is well-circumscribed and either nodular or smooth, as evident in the anterior segment photographs of both patients. It arises from the conjunctival stroma; therefore, the overlying epithelium is unchanged, which is one way of differentiating it from squamous-cell neoplasia.^{14,20}

Differential diagnoses for the clinical presentation of these lesions range from benign entities such as pinguecula, conjunctivitis and pterygium, to malignant ones such as Kaposi sarcoma, ocular surface squamous neoplasia, and amelanotic melanoma.^{4,20} It is important to refer any suspicious or atypical lesions or any non-resolving conjunctivitis for evaluation and biopsy, especially because clinical presentation alone cannot differentiate lymphoma from benign LH.

What additional tests/workup are indicated when ocular adnexal lymphoproliferative disease is suspected?

A thorough workup is needed to diagnose a lesion as lymphoproliferative disease. Careful staging of the disease is needed to determine whether the lesion is localized to the ocular adnexa or if there is concurrent systemic disease. Additional testing should include a physical examination with emphasis on the lymph nodes. Laboratory testing includes a CBC with differential, serum protein electrophoresis, serum lactate dehydrogenase and B_2 -microglobulin. Tissue biopsy with histological, immunophenotypic and molecular genetic evaluation is necessary, as are orbital and whole-body imaging.^{4,18} In some cases, a bone marrow biopsy may also be necessary if invasion of bone is suspected with more aggressive tumor types.⁶

While neither patient in the cases presented underwent magnetic resonance imaging (MRI) or computed tomography (CT), MRI and/or CT may aid in making the diagnosis as well as in assessing the exact location and extent of lesions. MRI and CT should be performed with contrast and thin slices through the orbit. On imaging studies, lymphoproliferative lesions typically appear as unifocal, well-circumscribed, homogenous masses that are either isodense or slightly hyperdense to muscle, and they mildly enhance with contrast. The lesions usually mold to solid structures, such as the globe and orbit. Bone destruction is always absent in LH and rare with lymphomas and usually signifies a more aggressive lymphoma, such as diffuse large-B-cell lymphoma.^{4,21}

Additional whole-body imaging is also necessary to look for systemic involvement and to stage the disease. Traditional imaging is performed using CT of the chest, abdomen and pelvis to look for any systemic lesions in lymph nodes, organs or other mucosal sites. Valenzuela and colleagues found that imaging with fluorine 18 deoxyglucose positron emission tomography (FDG PET) upstaged 66% of patients compared with traditional CT scan, which led to a change in the ultimate management of these patients.⁵ Similarly, a study by English et al. demonstrated that FDG PET detected systemic lesions in 31% more cases than CT imaging.²² Conversely, FDG PET is less sensitive at detecting orbital lesions compared to CT (27-79% vs. 73-97%, respectively).^{5,22} It has been suggested that combined PET/CT imaging could enhance the detection of the location and extent of disease and ultimately aid in management and final outcome.⁵

Ocular adnexal lymphoproliferative disorders are clinically and radiologically indistinguishable for the most part, and a full pathological evaluation including histology, immunophenotyping and molecular genetics is needed to confirm the exact type of lymphoid proliferation and to distinguish between benign and malignant disease.¹² Morphologically, LH is seen as a dense infiltration of small B and T lymphocytes organized into well-defined reactive follicles.²³ T cells are usually found to be the predominant cell type with immunohistochemistry, which differs from B-cell lymphomas — the majority of OALD. The reactive germinal centers contain “tingible body” macrophages filled with cellular debris, while Dutcher bodies are absent, differentiating LH from marginal-zone lymphomas such as MALT

lymphoma.^{14,23} The presence of an infiltrate of small, mostly B-cell lymphocytes surrounding reactive follicles also characterizes MALT lymphoma (distinguishing it by definition from diffuse large-B-cell lymphoma) although a few large cells may be present as well.²³ Immunophenotyping may also aid in differentiating it from other malignant lymphomas. The cells usually express the B-cell antigens CD19, CD20 and CD22 but are negative for CD5, CD10 and CD23, while follicular lymphoma is positive for CD10 and mantle-cell lymphoma is positive for CD5.¹⁶ Flow cytometry in the first case presentation showed CD19, CD20 and CD22 cell markers, which is consistent with a diagnosis of MALT lymphoma, while the second case presentation revealed CD10 cells consistent with a diagnosis of benign RLH.

What is the pathogenesis of lymphoproliferative disease?

Lymphoproliferative disease likely arises initially as an immune response to an antigen as the tissue architecture resembles that of a stimulated lymph node.¹⁴ Both LH and MALT lymphoma occur in locations where lymphocytes are not normally located but converge at the site in response to an antigen. Such stimuli are either from chronic infection, which has been shown with *Helicobacter pylori* in gastric tumors, or an autoimmune disease, such as Hashimoto's thyroiditis in thyroid tumors.^{4,6} Infection with *Chlamydia psittaci* has also been proposed as a precipitating factor in OALD, but studies have shown wide geographic variability, including variability within the same region, and there is no consensus.^{4,24}

In the case of LH, it has been hypothesized that T-cell imbalance drives B-cell proliferation because T cells are often the main cell type in LH and the CD4:CD8 ratio is usually elevated.²³ Also, it has been shown that LH can progress to NHL in cases of helicobacter gastritis, Sjögrens disease and Hashimoto's disease.²³ It is hypothesized that sustained B-cell proliferation in LH leads to increased mutations and translocation of genes until there is unrestrained proliferation of a monoclonal B-cell population (essentially the definition MALT lymphoma). However, not all lymphoid proliferations progress to lymphoma.²³

The exact pathogenesis of OALD is still unknown and investigation into the precipitating factors and the genetic mutations involved are ongoing.

What is the treatment for ocular adnexal lymphoproliferative disease?

The management of OALD involves multiple specialties and depends on tumor classification, extent of disease and prognostic indicators.^{2,4} It should only be decided after a rigorous examination, pathological evaluation and staging process.

The traditional treatment for primary OAL is radiotherapy, with a mean total dose of 32 Gy (range 15-46).²⁵ The patient in our first case report received a total of 36 Gy. The 5-year survival rate can be as high as 95-100% for orbital MALT lymphoma; however, higher doses of radiation or adjuvant treatment are usually needed in higher-grade tumors.¹⁹ Radiotherapy has also been used for LH, but at lower levels, typically 15-25 Gy. Standard treatments in the past have included systemic or injected steroids.¹⁴ Other treatments for OALD include observation only, which may be more appropriate in cases of LH as demonstrated in the second case report, and is not recommended for lymphoma unless the patient is particularly elderly and frail.²⁶ Surgical excision of encapsulated lesions is another treatment option, although excision has a high rate of recurrence and risk of possible systemic dissemination, especially if no adjuvant radiation or systemic treatment is added.^{4,11,26}

Adverse effects from radiotherapy can be categorized as acute (either during or within 3 months of treatment) or late (more than 3 months after treatment) and can occur in up to 70% of patients.^{19,26} Acute complications are mostly transient and often resolve spontaneously.^{25,27} Late effects are often chronic and more severe and may include dry eye syndrome, loss of eyelashes, cataracts, retinopathy and optic neuropathy. Radiation retinopathy typically occurs with doses of 45-60 Gy but may occur with doses as

low as 18 Gy when there is underlying vascular compromise such as diabetes.

For secondary orbital lymphoma with systemic involvement, systemic chemotherapy is typically the first-line treatment. Treatment of low-grade lymphoma often involves a single drug such as chlorambucil, while a multiagent regimen such as CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone) is used for more aggressive histological types.¹⁰ Combination treatment with both local radiation and systemic chemotherapy, termed chemoradiotherapy, may offer more favorable outcomes in patients with more advanced disease, but there have been no large trials confirming the optimal protocols.²⁸ Complications from systemic chemotherapy include nausea, anemia and neutropenia, which may be life-threatening.²⁹

In a small, retrospective study by Paik et al. in 2012, researchers compared 24 patients with non-conjunctival ocular adnexal MALT lymphoma.²⁸ Nine patients received chemotherapy only, eight received radiotherapy only (30-40 Gy), and seven received adjuvant radiotherapy (30-36 Gy) after chemotherapy provided insufficient response. They scored the ophthalmologic outcomes based on six main criteria that they deemed most common and important: decreased vision, dry eye, cataract, intractable intraocular pressure, retinopathy and blepharitis. They found that visual outcomes, as well as dry eye and blepharitis, were more common and serious in patients who received either radiotherapy alone or chemoradiotherapy vs. chemotherapy alone. Overall, patients treated solely with chemotherapy had better ocular and visual outcomes, and it should be considered as an alternative to radiotherapy.²⁹ Other case reports have also shown superior ophthalmologic outcomes with chemotherapy for primary disease. Dimitrakopoulos and colleagues stated that “although radiotherapy is preferable for localized lymphoproliferative lesions, chemotherapy also should be considered as an effective treatment that preserves the integrity and function of the ocular adnexa.”²⁹

Immunotherapy, most notably using the monoclonal antibody rituximab, is an emerging treatment for OALD. Rituximab targets the CD20 cell-surface receptor, which is found on all normal B cells and the majority of malignant B cells, leading to apoptosis. Single-agent systemic treatment with rituximab for patients who have received no prior treatment shows overall response rates of 50-100%, with either partial or complete remission.^{4,26} Unfortunately short-term recurrence is common, although a small case series by Annibali et al. showed improved results with maintenance therapy.^{4,10,26} Therefore, despite its better toxicity profile, rituximab is usually not used as monotherapy but rather is often added to chemotherapy, which may reduce the risk of failure.^{9,25} Rituximab is also used to treat LH when steroids or radiotherapy has failed. Cases of complete remission have been reported after use of rituximab as a first-line treatment.^{10,14}

Radioimmunotherapy is another new treatment option that adds a radioactive isotope to an anti-CD20 antibody. The radiation is emitted over an area greater than 100 cell diameters. Nearby tumor cells that are CD20-negative are also affected, a process called the ‘crossfire effect’.^{10,14} Response rates are as high as 90%, but studies of long-term outcomes are still needed.¹⁰

What is the prognosis for ocular adnexal lymphoproliferative disease?

The prognosis of OALD ultimately depends on type of lesion, extent of disease and treatment modality. Studies have shown that 31-44% of patients with OAL also have systemic lymphoma at the time of diagnosis. Systemic involvement is more likely in patients with bilateral disease and in patients diagnosed with high-grade tumor types.^{11-20,20} Based on the Kaplan-Meier survival analysis, studies have shown that systemic disease is likely to occur in 7-8% of patients at 1 year, 15-17% at 5 years, and 28-33% at 10 years with a greater risk in patients with bilateral involvement.^{11,20} A study by McKelvie et al. found that lymphoma-related deaths occurred in 18% of patients with OAL over 5 years of follow-up, but only in 2% of patients with MALT.¹⁵ Mortality was higher for follicular lymphoma (33%) and diffuse large-cell lymphoma (38%), but was 100% for the more aggressive types, such as mantle-cell

lymphoma, peripheral T-cell lymphoma and natural killer-cell lymphoma.¹²

In general, the risk of systemic involvement and death is greater in the more aggressive histological types of lymphoma and in patients with a more advanced stage of disease. MALT lymphoma usually has an indolent course with low likelihood of systemic involvement and lymphoma-related death regardless of the treatment modality selected. However, due to the risk of metastasis and possible transformation to a more aggressive lymphoma, all patients with OAL should be followed every 6 months for at least 5 years, including physical exam, laboratory tests and imaging.¹¹

Conclusion

We presented two cases of OALD that initially appeared similar on clinical examination, but after further evaluation with biopsy, immunohistochemistry, and molecular genetics were found to be respectively malignant and benign.

Advances in diagnostic techniques have allowed for better differentiation and classification of OALD. However, benign and malignant OALD are still indistinguishable on clinical examination and further testing is required. New MRI protocols may begin to be able to differentiate between benign LH and malignant lymphoma, but a histomorphological and immunohistochemical evaluation is still needed to confirm the exact histological type, which ultimately guides management. Whole-body imaging is needed to discover any systemic involvement, and combined PET/CT might prove to be the most sensitive way to simultaneously detect both orbital and systemic disease.

In addition to the traditional treatments of radiotherapy and chemotherapy for primary and systemic OALD, respectively, emerging treatments might offer better outcomes with less adverse effects. Immunotherapy with rituximab and radioimmunotherapy show promising results, especially when added as an adjunct to traditional treatments. New insights into the underlying etiology of lymphoproliferative disorders might also shed light on new treatments such as antibiotics, but there is no conclusive evidence at this time.

Benign lymphoid hyperplasia and MALT lymphoma of the ocular adnexa have an indolent course, and systemic involvement and tumor-related deaths are rare given their favorable response to treatment. However, progression and metastasis are risks, and patients should be monitored closely for at least 5 years, even after remission has occurred.

References

1. Demirci H, Shields CL, Karatza EC, Shields JA. Orbital lymphoproliferative tumors: analysis of clinical features and systemic involvement in 160 cases. *Ophthalmology*. 2008;115(9):1626-31, 1631.e1-3.
2. Sjö LD, Ralfkiaer E, Prause JU, et al. Increasing incidence of ophthalmic lymphoma in Denmark from 1980 to 2005. *Invest Ophthalmol Vis Sci*. 2008;49(8):3283-8.
3. Coupland SE, Krause L, Delecluse HJ, et al. Lymphoproliferative lesions of the ocular adnexa. Analysis of 112 cases. *Ophthalmology*. 1998;105(8):1430-41.
4. Valenzuela AA, Allen C, Grimes D, Wong D, Sullivan TJ. Positron emission tomography in the detection and staging of ocular adnexal lymphoproliferative disease. *Ophthalmology*. 2006;113(12):2331-7.
5. Sullivan TJ, Whitehead K, Williamson R, et al. Lymphoproliferative disease of the ocular adnexa: a clinical and pathologic study with statistical analysis of 69 patients. *Ophthal Plast Reconstr Surg*. 2005;21(3):177-88.
6. Coupland SE, Hummel M, Stein H. Ocular adnexal lymphomas: five case presentations and a review of the literature. *Surv Ophthalmol*. 2002;47(5):470-90.

7. Shields JA, Shields CL, Scartozzi R. Survey of 1264 patients with orbital tumors and simulating lesions: The 2002 Montgomery Lecture, part 1. *Ophthalmology*. 2004;111(5):997-1008.
8. Shinder R, Al-zubidi N, Esmaeli B. Survey of orbital tumors at a comprehensive cancer center in the United States. *Head Neck*. 2011;33(5):610-4.
9. Shome D, Esmaeli B. Targeted monoclonal antibody therapy and radioimmunotherapy for lymphoproliferative disorders of the ocular adnexa. *Curr Opin Ophthalmol*. 2008;19(5):414-21.
10. Alkatan HM, Alaraj A, El-khani A, Al-sheikh O. Ocular adnexal lymphoproliferative disorders in an ophthalmic referral center in Saudi Arabia. *Saudi J Ophthalmol*. 2013;27(3):227-30.
11. Oh DE, Kim YD. Lymphoproliferative diseases of the ocular adnexa in Korea. *Arch Ophthalmol*. 2007;125(12):1668-73.
12. Li EY, Yuen HK, Cheuk W. Lymphoproliferative disease of the orbit. *Asia Pac J Ophthalmol (Phila)*. 2015 Mar-Apr;4(2):106-11.
13. Moslehi R, Devesa SS, Schairer C, Fraumeni JF. Rapidly increasing incidence of ocular non-hodgkin lymphoma. *J Natl Cancer Inst*. 2006;98(13):936-9.
14. Verdijk RM. Lymphoproliferative Tumors of the Ocular Adnexa. *Asia Pac J Ophthalmol (Phila)*. 2017;6(2):132-142.
15. McKelvie PA. Ocular adnexal lymphomas: a review. *Adv Anat Pathol*. 2010;17(4):251-61.
16. Stefanovic A, Lossos IS. Extranodal marginal zone lymphoma of the ocular adnexa. *Blood*. 2009;114(3):501-10.
17. Bernardini FP, Bazzan M. Lymphoproliferative disease of the orbit. *Curr Opin Ophthalmol*. 2007 Sep;18(5):398-401.
18. Tovilla-canales JL, Tovilla y pomar JL, Ceron JR. Lymphoproliferative disorders of the ocular adnexa. *Curr Opin Ophthalmol*. 2004;15(5):401-5.
19. Shields CL, Chien JL, Surakiatchanukul T, Sioufi K, Lally SE, Shields JA. Conjunctival tumors: review of clinical features, risks, biomarkers, and outcomes ? The 2017 J. Donald M. Gass Lecture. *Asia Pac J Ophthalmol (Phila)*. 2017;6(2):109-120.
20. Sullivan TJ, Valenzuela AA. Imaging features of ocular adnexal lymphoproliferative disease. *Eye (Lond)*. 2006;20(10):1189-95.
21. Xu XQ, Hu H, Liu H, et al. Benign and malignant orbital lymphoproliferative disorders: Differentiating using multiparametric MRI at 3.0T. *J Magn Reson Imaging*. 2017;45(1):167-176.
22. Husain A, Roberts D, Pro B, McLaughlin P, Esmaeli B. Meta-analyses of the association between Chlamydia psittaci and ocular adnexal lymphoma and the response of ocular adnexal lymphoma to antibiotics. *Cancer*. 2007;110(4):809-15.
23. Andrew NH, Coupland SE, Pirbhai A, Selva D. Lymphoid hyperplasia of the orbit and ocular adnexa. A clinical pathologic review. *Surv Ophthalmol*. 2016;61(6):778-790
24. Finger PT. Radiation therapy for orbital tumors: concepts, current use, and ophthalmic radiation side effects. *Surv Ophthalmol*. 2009;54(5):545-68.
25. Annibali O, Chiodi F, Sarlo C, et al. Rituximab as Single Agent in Primary MALT Lymphoma of the Ocular Adnexa. *Biomed Res Int*. 2015;2015:895105.
26. Dhakal B, Fenske TS, Ramalingam S, et al. Local Disease Control in Ocular Adnexal Lymphoproliferative Disorders: Comparative Outcomes of MALT Versus Non-MALT Histologies. *Clin Lymphoma Myeloma Leuk*. 2017;17(5):305-311.e2.
27. Lim SH, Kang M, Son J. Extranodal marginal zone B cell lymphoma of mucosa-associated lymphoid tissue type of the ocular adnexa: retrospective single institution review of 95 patients. *Indian J Ophthalmol*. 2011 Jul-Aug;59(4):273-7.
28. Paik JS, Cho WK, Lee SE, et al. Ophthalmologic outcomes after chemotherapy and/or radiotherapy in non-conjunctival ocular adnexal MALT lymphoma. *Ann Hematol*. 2012;91(9):1393-401.
29. Dimitrakopoulos I, Venetis G, Kaloutsi V, Sotiropoulos D, Papadimitriou C. Effect of chemotherapy on primary mucosa-associated lymphoid tissue lymphoma of the orbit. *J Oral Maxillofac Surg*. 2008;66(1):16-20.

Dr. McCrary is a 2016 graduate of the State University of New York College of Optometry. She completed her residency in geriatrics/primary care optometry at the West Los Angeles Veterans Affairs Medical Center in 2017 and is currently employed by Kaiser Permanente in Bakersfield, Calif.

Dr. Yiasemis [andrea.yiasemis@va.gov] is an Assistant Professor at the Southern California College of Optometry at Marshall B. Ketchum University. She is a 2007 graduate of the University of California-Berkeley School of Optometry and she completed her residency at the San Francisco Veterans Affairs Medical Center. Her work experience includes San Francisco General Hospital and Veterans Affairs medical centers in Fresno and West Los Angeles, Calif.

Dr. Ilsen is a Professor at the Southern California College of Optometry (SCCO) at Marshall B. Ketchum University and has served as the residency program coordinator at the West Los Angeles Veterans Affairs Medical Center since 1996. She graduated from SCCO in 1988 and completed a residency in hospital-based optometry at the West Los Angeles Veterans Affairs Medical Center.

Features

Editorial

The Use of Social Media to Enhance Academic Careers

Aurora Denial OD, FAAO | Optometric Education: Volume 45, Number 1 (Fall 2019)



Aurora Denial, OD, FAAO

In June 2019, I had the opportunity to attend a conference, “Writing, Publishing, and Social Media for Healthcare Professionals,” sponsored by Harvard Medical School and the Massachusetts Psychological Association. The conference was interesting and educational. The most enlightening aspect was information presented on the use of social media to enhance health professionals’ careers. I like to think of myself as a moderately technologically savvy person; however, when it comes to social media, I am a digital baby, a dedicated Facebook stalker, who rarely posts any information. Nevertheless, the topic captured my attention because I recognized the enormous impact of social media in dissemination of scholarship, collaboration and networking.

Why Social Media?

Social media allows impact and visibility through exposure to a large audience, control of your message and a stage to advocate for your prospective. As a Department Chair, I am often in the position of mentoring and guiding faculty who are seeking promotion and tenure at various points in their careers. The achievement of promotion and tenure usually involves the demonstration of excellence and impact in the areas of teaching, service and scholarship. Traditionally, scholarly activities for promotion involve the production of high-quality scholarship and sharing the work via dissemination in peer-reviewed journals and presentations at national meetings. These activities eventually result in opportunities for national recognition and the development of a reputation for high-quality, impactful work in a particular area. Although faculty still need to follow this road map, the message from the conference was clear: In today’s world the traditional path alone is not sufficient.

Elsevier, a major participant in publishing and information analytics, reports that “more than 2.5 million scientific articles are published each year,” and “it has never been more important to ensure that your article stands out.”¹ Faculty interested in promotion and tenure need to take control to ensure their message gets out in a timely manner to an appropriate audience. Social media allows for the design, distribution and collaboration of material that can be measured.²

Social media includes social networking sites, which are defined as “web-based services that allow individuals to (1) construct a public or semi-public profile within a bounded system, (2) articulate a list of other users with whom they share a connection, and (3) view and traverse their list of connections and those made by others within the system.”³ Social networking can be earned (“exposure obtained by earning your way onto another’s platform”), rented (“a presence or content you control but lives on someone’s else’s platform”) or owned (“any channel where you fully own the connection to your audience”).⁴ Owned media include your website, blog and e-mail list, assuming they live in a domain that you own. Owned media provides an opportunity to develop your own audience. Rented media includes Facebook, Twitter, LinkedIn, Instagram etc. These sites provide opportunities to collaborate, network and post your latest accomplishments, which can include research findings, articles, images, videos and audio recordings. Earned media such as an interview on a podcast gives you instant access to an

existing audience. However, the downside is that you need to be invited. Academic careers are impacted when faculty understand and utilize all three types of networks.⁴ According to Elsevier, “every day scholarly articles receive thousands of new mentions across social media, news and blogs.”¹

Where to Start

Social media presents opportunities as well as challenges for academicians. It can seem overwhelming to learn and navigate as well as time-consuming to maintain. That said, many institutions have dedicated staff to help faculty efficiently and effectively use social media. Elsevier has provided extensive information on what you can do to get your article the attention it deserves.¹ Embracing social media as a means of career enhancement may seem foreign and disruptive to many. Using more traditional methods of building a career may initially seem more comfortable. However, faculty and mentors need to be open-minded and aware of the current landscape. Opportunities that arise from social media use may take years to achieve by more traditional means. It is a disservice to the faculty members we mentor to not explore all avenues that may be beneficial to their careers.

References

1. Get noticed. Promoting your article for maximum interest. [Internet]. Elsevier; c2005 [cited 2019 Oct 10]. Available from: https://www.elsevier.com/_data/assets/pdf_file/0013/201325/Get-Noticed_Brochure_2018.pdf
2. Cabrera D, Vartabedian BS, Spinner RJ, et al. More than likes and tweets: creating social media portfolios for academic promotion and tenure. *J Grad Med Educ.* 2017;9(4):421-425.
3. Boyd DM, Ellison NB. Social network sites: definition, history, and scholarship. *J Comput Mediat Commun.* 2007;13(1):210-230.
4. Hendricks B, Shelton R. Mastering the new media landscape. Embrace the micromedia mindset. Oakland, CA: Berrett-Koehler Publishers; 2016. Chapter 1.

Dr. Denial [deniala@neco.edu], Editor of *Optometric Education*, is a Professor and Chair of the Department of Primary Care at the New England College of Optometry and a Clinical Instructor at a community health center in Boston.

Industry News

Announcement

ASCO's Public Awareness Campaign "Optometry Gives Me Life" Surpasses Benchmarks

Keshia Elder, OD, MS, FAAO | *Optometric Education: Volume 45, Number 1 (Fall 2019)*

In the eight months since its March 2019 launch, ASCO's public awareness campaign "Optometry Gives Me Life" has gained impressive momentum. The online, e-mail and social media ads associated with the campaign have been viewed nearly 35 million times.¹ According to the media and communications agency partnered with ASCO on the initiative, the percentage of people who see the ads and click on them, known as the "click-through rate," is 80% higher than the benchmark typically achieved with this type of campaign. "Optometry Gives Me Life" also surpassed the benchmark for the percentage of people who watch campaign videos in their entirety. At 85.41%, the campaign's rate is 187% higher than the 30% benchmark.¹ In addition, nearly 99,000 people visited the campaign landing page, <https://futureeyedoc.org/>,¹ in the initial eight-months.

Why Doesn't Everyone See the Campaign Ads?

"Optometry Gives Me Life" is a multipronged and highly targeted campaign. Its intended audience is college juniors and seniors pursuing STEM degrees who have expressed an interest in a health profession. Internet users whose age and online activity don't match the target audience criteria won't see the ads. The ads present to those who fit the criteria and direct them to the <https://futureeyedoc.org/> landing page. There, they see the campaign videos and can access additional information.

Also as part of the campaign, each ASCO member school and college of optometry received a toolkit containing "Optometry Gives Me Life" materials they can customize and use to promote optometry, recruit optometry students and expand the reach of the campaign.

The Impetus for "Optometry Gives Me Life"

The "Optometry Gives Me Life" campaign supports one of ASCO's key strategic objectives, which is to attract a robust, highly qualified and diverse pool of applicants pursuing the education necessary for providing optometric vision care, thus improving the overall health of patients everywhere. The campaign addresses the concern that a recent decrease in the number of optometry school applicants and the resulting increase in seat availability may lead to less competition for optometry school acceptance and a less-qualified applicant pool. The number of optometry school applicants has been on the decline since 2015-2016,² and the applicant-to-matriculant ratio decreased from 1.53 (2503:1639) in 2010 to 1.39 (2527:1822) in 2018.³

[The Applicant Pool Development Task Force \(APDTF\)](#) is the steward of ASCO's applicant development efforts. Formed in 2017, it includes representatives from schools and colleges of optometry with expertise in admissions, marketing and communications; industry partners; and a representative from the ASCO Diversity and Cultural Competency Committee. In the Association's [Strategic Plan 2018-2022](#), applicant pool development is tied to measurable results:

- recruit at least two applicants with an undergraduate grade-point average higher than 3.0 for every available training slot in a United States-based optometry program by June 2022
- increase by 100% the number of highly qualified applicants from Hispanic/Latino and African American/Black backgrounds by June 2022
- maximize the percentage/number of high school students who are aware of optometry as a career

option

Off to a Good Start

It's too early to determine the overall impact *Optometry Gives Me Life* will have, but the early indicators are promising. By promoting and renewing interest in optometry as a career, the campaign is moving ASCO and its member institutions closer to the important goal of increasing the quantity, quality and diversity of optometry school applicants.

References

1. ASCO media campaign performance dashboard, "Optometry Gives Me Life" campaign. March 2019-present. Author: Butler/Till Media Services.
2. OptomCAS Applicant Data Reports [Internet]. Rockville, MD: Association of Schools and Colleges of Optometry [cited Oct 16, 2019]. Available from: <https://www.optomcas.org/>
3. Zadnik K, Reich LN. An analysis of the optometric applicant pool relative to matriculants. *Optom Vis Sci.* 2019 Sep;96(9):637-646.

Dr. Elder is an Associate Professor and Director of the externship program at the University of Alabama at Birmingham School of Optometry. She is an Associate Editor and peer reviewer for *Optometric Education* and works with ASCO as the Applicant Pool Development Task Force liaison to the Diversity and Cultural Competency Committee. Dr. Elder is also the Chair-Elect for the ASCO Diversity, Equity and Inclusion Special Interest Group

Industry News

Announcement

Two Join ASCO Journal as Associate Editors

Desiree Ifft | Optometric Education: Volume 45, Number 1 (Fall 2019)

Jamie Althoff, OD, and Keshia S. Elder, OD, MS, FAAO, have been selected as Associate Editors for ASCO's journal, *Optometric Education*.

Dr. Althoff, a 2007 graduate of the Ferris State University Michigan College of Optometry, is currently an Assistant Professor at the Nova Southeastern University College of Optometry where she also serves as the Director of Outcomes Assessment and Faculty Development. She teaches theoretical optics, clinical gerontology, and optometric theory and methods, and her academic interests include the scholarship of teaching and learning and the use of technology in education. Dr. Elder, a 1998 graduate of the University of Alabama at Birmingham School of Optometry (UAB), is currently an Associate Professor at UAB where she is also Director of the externship program. She precepts students in the Contact Lens and Primary Care clinics and is the course director for Clinical Management. Dr. Elder is a member of the American Academy of Optometry's Lectures and Workshops Committee and works with ASCO as the Applicant Pool Development Task Force liaison to the Diversity and Cultural Competency Committee. She is also Chair-Elect for the ASCO Diversity, Equity and Inclusion Special Interest Group.



Jamie Althoff, OD Keshia S. Elder, OD, MS, FAAO

Both Dr. Althoff and Dr. Elder are long-time members of the *Optometric Education* Journal Review Board. Dr. Elder says she wants to be more involved with the journal because it has the ability to positively impact the entire field of optometric education. "The publication is important because it's the only journal focused solely on optometric education," she says. "It's a place to go to learn about innovative teaching methods, brush up on clinical knowledge by reading the teaching case reports, and stay abreast of scholarship in optometric teaching and learning."

Optometric Education Editor Aurora Denial, OD, FAAO, says Drs. Elder and Althoff will be involved with implementing new projects, collaborating in selection of Journal Review Board members, soliciting manuscripts and assisting with the development of journal content. "I'm honored to have the opportunity to continue the hard work of those who have developed the journal into what it is today," Dr. Althoff says. "My colleagues and I are always striving to find the best way to teach and motivate the future optometrists in our programs, but most of us have little or no formal training in education. *Optometric Education* serves as a valuable resource for us to share our experiences and inspire each other. I hope to help it continue to adapt to meet the ever-changing needs of optometry and optometric education."

Industry News

